

REC'D 19 MAY 2004

WIPO PCT

PA 1137470

**THE UNITED STATES OF AMERICA****TO ALL TO WHOM THESE PRESENTS SHALL COME:****UNITED STATES DEPARTMENT OF COMMERCE****United States Patent and Trademark Office****March 03, 2004**


**THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE UNDER 35 USC 111.**

**APPLICATION NUMBER: 60/466,542****FILING DATE: April 30, 2003**

RULE 17.1(a) OR (b)

**PRIORITY DOCUMENT**  
SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH  
RULE 17.1(a) OR (b)

**By Authority of the**  
**COMMISSIONER OF PATENTS AND TRADEMARKS**

**E. BORNETT****Certifying Officer****BEST AVAILABLE COPY**

05-01-03 04466542 . 043003 / PROV 7  
SUBSTITUTE for Provisional Application for Patent Cover Sheet PTO/SB/16 (10-01)  
Approved for use through 10/31/2002. OMB 0651-0032  
U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

# PROVISIONAL APPLICATION FOR PATENT COVER SHEET

This is a request for filing a PROVISIONAL APPLICATION FOR PATENT under 37 CFR 1.53 (c).

DOCKET NUMBER

MC054PV

## INVENTOR(S)

Given Name (first and middle [if any])	Family Name or Surname	Residence (City and either State or Foreign Country)
Daniel Laurence Michel Patrick Denis Dwight	Dube Dube Gallant Lacombe Deschenes Macdonald	1811 Blueberry Forest, St-Lazare, Quebec, Canada J7T 2J9 16729 Blvd. de Pierrefonds, App. 301, Pierrefonds, Quebec, Canada H9H 5L8 15 Ballantyne South, Montreal, Quebec, Canada H4X 2B1 139 Villeneuve Ouest, Montreal, Quebec, Canada 43T-2R6 1445 Carson Dorval, Quebec, Canada H9S 1M7 3216 Cherrier, L'Ile Bizard, Quebec, Canada H9C 1E2

☐ Additional inventors are being named on the separately numbered sheets attached hereto

## TITLE OF THE INVENTION (500 characters max)

8-(3-BIARYL)PHENYYQUINOLINE PHOSPHODIESTERASE-4 INHIBITORS

## CORRESPONDENCE ADDRESS

Direct all Correspondence to:

Merck & Co., Inc.  
Patent Department - RY60-30  
P.O. Box 2000  
Rahway

☒ Customer Number

000210

STATE

New Jersey

ZIP CODE

07065

COUNTRY

U.S.A.

## ENCLOSED APPLICATION PARTS (check all that apply)

☒ Specification

Number of Pages

126

☐ CD(s), Number

☐ Drawing(s)

Number of Sheets

☐ Other (specify)

☐ Application Data Sheet. See 37 CFR 1.76

## METHOD OF PAYMENT OF FILING FEES FOR THIS PROVISIONAL APPLICATION FOR PATENT (check one)

☐ A check or money order is enclosed to cover the filing fees

☒ The Commissioner is hereby authorized to charge filing fees or credit any overpayment to Deposit Account Number:

13-2755

FILING FEE  
AMOUNT (\$)

\$160.00

The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.

☒ No.

☐ Yes, the name of the U.S. Government agency and the Government contract number are:

Respectfully submitted,

SIGNATURE

*Curtis C. Panzer*

Date 04/30/2003

TYPED or PRINTED NAME Curtis C. Panzer

REGISTRATION NO. 33,752  
(if appropriate)

TELEPHONE 732-594-3199

## EXPRESS MAIL CERTIFICATE

DATE OF DEPOSIT April 30, 2003

EXPRESS MAIL NO. EL920632540US

I HEREBY CERTIFY THAT THIS CORRESPONDENCE IS BEING DEPOSITED WITH THE UNITED STATES POSTAL SERVICE AS EXPRESS MAIL "POST OFFICE TO ADDRESSEE" ON THE ABOVE DATE IN AN ENVELOPE ADDRESSED TO ASSISTANT COMMISSIONER FOR PATENTS, WASHINGTON, D.C. 20231

MAILED BY

*Kelly J. Pano*

DATE

4-30-2003

## TITLE OF THE INVENTION

8-(3-BIARYL)PHENYLQUINOLINE PHOSPHODIESTERASE-4 INHIBITORS

## BACKGROUND OF THE INVENTION

## FIELD OF THE INVENTION

5 The present invention is directed to compounds that are substituted 8-phenylquinolines. In particular, this invention is directed to substituted 8-phenylquinolines which are phosphodiesterase-4 inhibitors wherein the phenyl group  
10 at the 8-position contains a *meta* aryl or heterocyclic group.

## RELATED BACKGROUND

Hormones are compounds that variously affect cellular activity. In many respects, hormones act as messengers to trigger specific cellular responses and  
15 activities. Many effects produced by hormones, however, are not caused by the singular effect of just the hormone. Instead, the hormone first binds to a receptor, thereby triggering the release of a second compound that goes on to affect the cellular activity. In this scenario, the hormone is known as the first messenger while the second compound is called the second messenger. Cyclic adenosine monophosphate  
20 (adenosine 3', 5'-cyclic monophosphate, "cAMP" or "cyclic AMP") is known as a second messenger for hormones including epinephrine, glucagon, calcitonin, corticotrophin, lipotropin, luteinizing hormone, norepinephrine, parathyroid hormone, thyroid-stimulating hormone, and vasopressin. Thus, cAMP mediates cellular responses to hormones. Cyclic AMP also mediates cellular responses to various  
25 neurotransmitters.

Phosphodiesterases ("PDE") are a family of enzymes that metabolize 3', 5' cyclic nucleotides to 5' nucleoside monophosphates, thereby terminating cAMP second messenger activity. A particular phosphodiesterase, phosphodiesterase-4 ("PDE4", also known as "PDE-IV"), which is a high affinity, cAMP specific, type IV  
30 PDE, has generated interest as potential targets for the development of novel anti-asthmatic and anti-inflammatory compounds. PDE4 is known to exist as at least four isoenzymes, each of which is encoded by a distinct gene. Each of the four known PDE4 gene products is believed to play varying roles in allergic and/or inflammatory responses. Thus, it is believed that inhibition of PDE4, particularly the specific PDE4  
35 isoforms that produce detrimental responses, can beneficially affect allergy and

inflammation symptoms. It would be desirable to provide novel compounds and compositions that inhibit PDE4 activity.

A major concern with the use of PDE4 inhibitors is the side effect of emesis which has been observed for several candidate compounds as described in  
5 C.Burnouf et al., ("Burnouf"), *Ann. Rep. In Med. Chem.*, 33:91-109(1998). B.Hughes et al., *Br. J.Pharmacol.*, 118:1183-1191(1996); M.J.Perry et al., *Cell Biochem. Biophys.*, 29:113-132(1998); S.B.Christensen et al., *J.Med. Chem.*, 41:821-835(1998); and Burnouf describe the wide variation of the severity of the undesirable side effects exhibited by various compounds. As described in M.D.Houslay et al., *Adv. In Pharmacol.*, 44:225-342(1998) and D.Spina et al., *Adv. In Pharmacol.*, 44:33-89(1998), there is great interest and research of therapeutic PDE4 inhibitors.

International Patent Publication WO9422852 describes quinolines as PDE4 inhibitors. U.S. Patent No. 6,410,563 describes 8-arylquinoline PDE4 inhibitors.

15 A.H.Cook, et al., *J.Chem. Soc.*, 413-417(1943) describes gamma-pyridylquinolines. Other quinoline compounds are described in Kei Manabe et al., *J.Org. Chem.*, 58(24):6692-6700(1993); Kei Manabe et al., *J.Am. Chem. Soc.*, 115(12):5324-5325(1993); and Kei Manabe et al., *J.Am. Chem. Soc.*, 114(17):6940-6941(1992).

20 Compounds that include ringed systems are described by various investigators as effective for a variety of therapies and utilities. For example, International Patent Publication No. WO 98/25883 describes ketobenzamides as calpain inhibitors, European Patent Publication No. EP 811610 and U.S. Patent Nos. 5,679,712, 5,693,672 and 5,747,541 describe substituted benzoylguanidine sodium  
25 channel blockers, U.S. Patent No. 5,736,297 describes ring systems useful as a photosensitive composition.

U.S. Patent Nos. 5,491,147, 5,608,070, 5,622,977, 5,739,144, 5,776,958, 5,780,477, 5,786,354, 5,798,373, 5,849,770, 5,859,034, 5,866,593, 5,891,896, and International Patent Publication WO 95/35283 describe PDE4  
30 inhibitors that are tri-substituted aryl or heteroaryl phenyl derivatives. U.S. Patent No. 5,580,888 describes PDE4 inhibitors that are styryl derivatives. U.S. Patent No. 5,550,137 describes PDE4 inhibitors that are phenylaminocarbonyl derivatives. U.S. Patent No. 5,340,827 describes PDE4 inhibitors that are phenylcarboxamide compounds. U.S. Patent No. 5,780,478 describes PDE4 inhibitors that are tetra-  
35 substituted phenyl derivatives. International Patent Publication WO 96/00215

MC054PV

describes substituted oxime derivatives useful as PDE4 inhibitors. U.S. Patent No. 5,633,257 describes PDE4 inhibitors that are cyclo(alkyl and alkenyl)phenyl-alkenyl (aryl and heteroaryl) compounds.

5 However, there remains a need for novel compounds and compositions that therapeutically inhibit PDE4 with minimal side effects.


#### SUMMARY OF THE INVENTION

10 The present invention is directed to novel substituted 8-phenylquinolines that are PDE4 inhibitors, wherein the phenyl group at the 8-position is substituted by an aryl or heterocyclic group in the *meta* position. This invention also provides a pharmaceutical composition which includes an effective amount of the novel substituted 8-phenylquinoline and a pharmaceutically acceptable carrier. This invention further provides a method of treatment in mammals of, for example, asthma, chronic bronchitis, chronic obstructive pulmonary disease (COPD),  
15 eosinophilic granuloma, psoriasis and other benign or malignant proliferative skin diseases, endotoxic shock (and associated conditions such as laminitis and colic in horses), septic shock, celiac sprue, ulcerative colitis, Crohn's disease, reperfusion injury of the myocardium and brain, inflammatory arthritis, chronic glomerulonephritis, atopic dermatitis, urticaria, adult respiratory distress syndrome,  
20 chronic obstructive pulmonary disease in animals, diabetes insipidus, allergic rhinitis, allergic conjunctivitis, vernal conjunctivitis, arterial restenosis, orthrosclerosis, atherosclerosis, neurogenic inflammation, pain, cough, rheumatoid arthritis, ankylosing spondylitis, transplant rejection and graft versus host disease, hypersecretion of gastric acid, bacterial, fungal or viral induced sepsis or septic shock,  
25 inflammation and cytokine-mediated chronic tissue degeneration, osteoarthritis, cancer, cachexia, muscle wasting, depression, memory impairment, tumour growth and cancerous invasion of normal tissues by the administration of an effective amount of the novel substituted 8-phenylquinoline.

#### 30 DETAILED DESCRIPTION OF THE INVENTION

In a first aspect the invention is directed to compounds represented by Formula (I):

MC054PV



(I)

and pharmaceutically acceptable salts, wherein

S<sub>1</sub>, S<sub>2</sub>, and S<sub>3</sub> are independently

1. H,
2. -OH,
3. halogen,
4. -C<sub>1</sub>-C<sub>6</sub>alkyl,
5. -O-C<sub>1</sub>-C<sub>6</sub>alkyl optionally substituted with 1, 2 or 3 halogens, or
6. -CN;

R<sub>1</sub> is

1. -(C<sub>1</sub>-C<sub>6</sub>alkyl)-SO<sub>n</sub>-(C<sub>1</sub>-C<sub>6</sub>alkyl) group, optionally substituted with 1, 2 or 3 substituents; wherein each substituent is independently a halogen, -OH and -CN,
2. -C(O)-O-aryl,
3. -C(O)-NH-aryl,
4. -C(O)-NH-heterocycle or N-oxide thereof,
5. -C(O)-NH-C<sub>1</sub>-C<sub>6</sub>alkyl,
6. -C(O)-NH-cycloC<sub>3</sub>-C<sub>6</sub>alkyl,
7. -C<sub>1</sub>-C<sub>6</sub>alkyl, optionally substituted with 1 to 6 halogens and 1 hydroxy,
8. -COOH,
9. -C<sub>1</sub>-C<sub>6</sub>alkyl-COOH,
10. -O-C<sub>1</sub>-C<sub>6</sub>alkyl,
11. -cycloC<sub>3</sub>-C<sub>6</sub>alkyl,
12. -C<sub>3</sub>-C<sub>6</sub>alkyl-heterocycle,
13. aryl,

MC054PV

14. heterocycle,
15. carbonyl,
16. carbamoyl, or
17.  $\text{-SO}_n\text{-(C}_1\text{-C}_6\text{alkyl)}$ ;

each  $n$  is independently 0, 1, or 2;

$\text{Ar}_1$  and  $\text{Ar}_2$  are each independently an aryl or heterocycle or an N-oxide thereof;

$\text{R}_2$  is

1. Hydrogen,
2. aryl optionally substituted with 1, 2 or 3 substituents selected from halogen,
3. heterocycle optionally substituted with 1, 2 or 3 halogens,
4.  $\text{-C}_1\text{-C}_6\text{alkyl}$  optionally substituted with 1, 2 or 3 substituents selected from hydroxy and halogen,
5.  $\text{-COOH}$ ,
6. 1, 2 or 3 halogens,
7.  $\text{-SO}_n\text{-(C}_1\text{-C}_6\text{alkyl)}$ ,
8.  $\text{-N(H)-S(O)}_n\text{-C}_1\text{-C}_6\text{alkyl}$ ,
9.  $\text{-O-C}_1\text{-C}_6\text{alkyl}$  substituents each optionally substituted with 1, 2 or 3 halogens,
10.  $\text{-C(O)-N(H)-C}_3\text{-C}_6\text{cycloalkyl}$ , or
11.  $\text{-C(O)-C}_1\text{-C}_6\text{alkyl}$ ;

$\text{R}_3$  is

1. Hydrogen,
2.  $\text{-C}_1\text{-C}_6\text{alkyl}$  optionally substituted with hydroxy,  $\text{-S(O)}_n\text{C}_1\text{-C}_6\text{alkyl}$ , heterocycle, or 1, 2, 3, 4, 5 or 6 halogens,
3. aryl or  $\text{C}_6\text{-C}_{12}\text{cycloalkyl}$  optionally substituted with phenyl,  $\text{-C}_1\text{-C}_6\text{alkyl}$ ,  $\text{-S(O)}_n\text{C}_1\text{-C}_6\text{alkyl}$ ,  $\text{-C(O)-O-C}_1\text{-C}_6\text{alkyl}$ ,  $\text{-COOH}$ , hydroxy- $\text{C}_1\text{-C}_6\text{alkyl}$ - or 1, 2 or 3 halogens,
4. heterocycle or optionally substituted with 1, 2 or 3 substituents independently selected from phenyl, halogen,  $\text{C}_1\text{-C}_6\text{alkyl}$ , hydroxy- $\text{C}_1\text{-C}_6\text{alkyl}$ ,  $\text{-COOH}$ ,  $\text{-C(O)-O-C}_1\text{-C}_6\text{alkyl}$ ,
5. amino,
6.  $\text{-C(O)-O-C}_1\text{-C}_6\text{alkyl}$ ,

MC054PV

7.  $-C_1-C_6\text{alkyl-O-phenyl}$  optionally substituted with 1, 2 or 3 halogens,
  8.  $-C_1-C_6\text{alkyl-phenyl}$  optionally substituted with 1 or 2 substituents selected from hydroxy and halo,
  9.  $-COOH$ ,
  10. Halogen,
  11.  $-SO_n-(C_1-C_6\text{alkyl})$ ,
  12.  $-N(H)-S(O)_n-C_1-C_6\text{alkyl}$  optionally substituted with 1, 2 or 3 halogen,
  13.  $-N(H)-C(O)-C_1-C_6\text{alkyl}$ ,
  14.  $-N(H)\text{-heterocycle}$  optionally substituted with 1, 2 or 3 halogens,
  15.  $-N(H)\text{-aryl}$  optionally substituted with 1, 2 or 3 halogens,
  16.  $-N(H)-C_1-C_6\text{alkyl}$  optionally substituted with 1, 2 or 3 halogens,
  17.  $-C(O)-N(H)-C_1-C_6\text{alkyl}$  optionally substituted with 1, 2 or 3 halogens,
  18.  $-C(O)-NH-C_3-C_6\text{cycloalkyl}$ ,
  19.  $-O-C_1-C_6\text{alkyl}$  optionally substituted with 1, 2 or 3 halogens or phenyl optionally substituted with 1, 2, or 3 halogen;
- $R_4$  is
1. H,
  2. Halogen,
  3.  $-CN$
  4.  $-C_1-C_6\text{alkyl}$ ,
  5.  $-O-C_1-C_6\text{alkyl}$  optionally substituted with 1, 2 or 3 halogens,
  6.  $-C_1-C_6\text{alkyl-phenyl}$  with phenyl optionally substituted with halogen, or
  7. Oxo.

Within this first aspect is the genus of compounds and pharmaceutically acceptable salts thereof, wherein  $Ar_1$  is pyridine or pyridinone or an N-oxide thereof.

Within this genus is the sub-genus of compounds and pharmaceutically acceptable salts thereof, wherein



MC054PV

Ar<sub>2</sub> is phenyl, oxadiazole or thiadiazole, tetrazole.

Within this sub-genus is the class of compounds and pharmaceutically acceptable salts thereof, wherein

5 R<sub>1</sub> is -(C<sub>1</sub>-C<sub>6</sub>alkyl)-SO<sub>n</sub>-(C<sub>1</sub>-C<sub>6</sub>alkyl); and  
R<sub>2</sub> is -SO<sub>n</sub>-C<sub>1</sub>-C<sub>6</sub>alkyl.

Also within this first aspect, there is a genus of compounds and pharmaceutically acceptable salts thereof, wherein

10 Ar<sub>1</sub> is phenyl.

Within this genus there is a sub-genus of compounds and pharmaceutically acceptable salts thereof, wherein

Ar<sub>2</sub> is phenyl, oxadiazole, thiadiazole, tetrazole, pyridine or pyridinone or an N-oxide thereof.

15 Within this sub-genus there is a class of compounds and pharmaceutically acceptable salt thereof, wherein

R<sub>1</sub> is -(C<sub>1</sub>-C<sub>6</sub>alkyl)-SO<sub>n</sub>-(C<sub>1</sub>-C<sub>6</sub>alkyl); and  
R<sub>2</sub> is -SO<sub>n</sub>-C<sub>1</sub>-C<sub>6</sub>alkyl.

20 Also within the first aspect there is a genus of compounds and pharmaceutically acceptable salts thereof, wherein

Ar<sub>1</sub> is thiazole or oxazole.

25 Within this genus there is a sub-genus of compounds and pharmaceutically acceptable salts thereof, wherein

Ar<sub>2</sub> is phenyl, pyridine or pyridinone or an N-oxide thereof.


30 Within this sub-genus of compounds there is a class of compounds and pharmaceutically acceptable salts thereof, wherein

R<sub>1</sub> is -(C<sub>1</sub>-C<sub>6</sub>alkyl)-SO<sub>n</sub>-(C<sub>1</sub>-C<sub>6</sub>alkyl); and  
R<sub>2</sub> is -SO<sub>n</sub>-C<sub>1</sub>-C<sub>6</sub>alkyl.

MC054PV

Also within the first aspect there is a genus of compounds of Formula

Ia



and a pharmaceutically acceptable salts thereof, wherein:

Ar<sub>1</sub> is phenyl, pyridine, pyridinone, pyrimidyl, thiophene, thiazole, triazole, tetrazole, oxazole, thienodiazole, pyridodiazole, imidazothiazole or quinoxaline or an N-oxide thereof; and

Ar<sub>2</sub> is phenyl, pyridine, pyridinone, oxadiazole, tetrazole or thiadiazole or an N-oxide thereof.

Within this genus of compounds of Formula Ia there is a sub-genus of compounds and pharmaceutically acceptable salts thereof, wherein:

R<sub>2</sub> is phenyl, -COOH, -C<sub>1</sub>-C<sub>6</sub>alkyl, -C<sub>1</sub>-C<sub>6</sub>alkoxy, mono or di-halo-C<sub>1</sub>-C<sub>6</sub>alkoxy, -hydroxyC<sub>1</sub>-C<sub>6</sub>alkyl, or -SO<sub>n</sub>-(C<sub>1</sub>-C<sub>6</sub>alkyl) or 1, 2 or 3 halogens.

Within this sub-genus there is a class of compounds of Formula Ia and pharmaceutically acceptable salt thereof, wherein:

R<sub>3</sub> is Hydrogen, amino, biphenyl, N-(tert-butoxycarbonyl)-4-phenylpyrrolidin-3-yl, N-(tert-butoxycarbonyl)azetidin-3-yl, N-(tert-butoxycarbonyl)pyrrolidin-3-yl, 3-chloro-4-fluorophenyl, 4-chlorophenoxymethyl, 2-chlorophenyl, 4-chlorophenyl, ethoxycarbonyl, furan-2-yl, furan-3-yl, imidazol-2-yl, indan-1-yl, indan-2-yl, 1H-indol-2-yl, 1H-indol-3-yl, 1H-indol-4-yl, 1H-indol-5-yl, 1H-indol-6-yl, 1H-indol-7-yl, isoquinolin-1-yl, isoquinolin-4-yl, isoquinolin-5-yl,

MC054PV

isoquinolin-8-yl, isoxazol-3-yl, 3-methoxycarbonylphenyl, 4-  
 methoxycarbonylphenyl, methyl, 1-methyl-1Hpyrazol-3-yl, 1-methyl-1Hpyrazol-4-yl,  
 1-methyl-1Hpyrazol-5-yl, 2-methylphenyl, 3-methylphenyl, 4-methylphenyl, 2-  
 methylpyridin-5-yl, methylsulfonylmethyl, 2-methylsulfonylphenyl, 3-  
 5 methylsulfonylphenyl,  
 4-methylsulfonylphenyl, morpholin-4-ylmethyl, phenyl, pyrazinyl, 1H-pyrazol-3-yl,  
 pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, pyridin-2-yl N-oxide, pyridin-3-yl N-oxide,  
 pyridin-4-yl N-oxide, 3-pyridinylmethyl, pyrimidin-2-yl, pyrimidin-4-yl, pyrimidin-5-  
 yl, quinolin-4-yl, quinolin-5-yl, quinolin-8-yl, 5,6,7,8-tetrahydro-1,8-naphthyridin-2-  
 10 yl, 5,6,7,8-tetrahydro-5H-benzo[a][7]annulen-5-yl, 5,6,7,8-tetrahydro-5H-  
 benzo[a][7]annulen-6-yl, tetrahydrofuran-2-yl, 1,2,3,4-tetrahydronaphthalen-1-yl,  
 1,2,3,4-tetrahydronaphthalen-2-yl, 1,3-thiazol-2-yl, 1,3-thiazol-5-yl, thiophen-2-yl and  
 thiophen-2-yl, 3-(propan-2-ol)-phenyl, 4-(propan-2-ol)-phenyl, propan-2-ol,  
 15 hexafluoropropan-2-ol.

Also within the genus of compounds of Formula Ia there is the sub-genus of compounds and pharmaceutically acceptable salts thereof, wherein:

Ar<sub>1</sub> is thiazole;

Ar<sub>2</sub> is phenyl; and

20 R<sub>2</sub> is -SO<sub>2</sub>-C<sub>1</sub>-C<sub>6</sub>alkyl or halogen or C<sub>1</sub>-C<sub>6</sub>alkyl optionally substituted with hydroxy or 1-3 halogens.

Within this sub-genus there is the class of compounds and pharmaceutically acceptable salts thereof, wherein:

25 R<sub>3</sub> is

1. Hydrogen,

2. -C<sub>1</sub>-C<sub>6</sub>alkyl optionally substituted with hydroxy, -S(O)<sub>n</sub>C<sub>1</sub>-C<sub>6</sub>alkyl, or 1-6 halogens.

30 For Example, R<sub>3</sub> may be propan-2-ol, hexafluoropropan-2-ol, such that R<sub>3</sub>-Ar may be 3-(propan-2-ol)-phenyl or 3-methylsulfonylphenyl.

Also within the genus of compounds of Formula Ia there is the sub-genus of compounds and pharmaceutically acceptable salts thereof, wherein

Ar<sub>1</sub> is pyridine or an N-oxide thereof;

35 Ar<sub>2</sub> is oxadiazole; and

MC054PV

R<sub>2</sub> is

1. -C<sub>1</sub>-C<sub>6</sub>alkyl optionally substituted with hydroxy, -S(O)<sub>n</sub>C<sub>1</sub>-C<sub>6</sub>alkyl, or 1-3 substituents halogens,
2. -N(H)-C(O)-C<sub>1</sub>-C<sub>6</sub>alkyl,
3. -COOH, or
4. -C(O)-NH-C<sub>3</sub>-C<sub>6</sub>cycloalkyl.

Within this sub-genus of compounds of Formula Ia there is a class of compounds and pharmaceutically and acceptable salts thereof wherein:

R<sub>3</sub> is hydrogen.

Illustrating the invention are:

- 8-(3-{2-(3-chlorophenyl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-5-yl}phenyl)-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline,
- 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-{3-[4-[4-(methylsulfonyl)phenyl]-2-(1-oxidopyridin-4-yl)-1,3-thiazol-5-yl]phenyl}quinoline,
- 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-{3-[4-[4-(methylsulfonyl)phenyl]-2-(1-oxidopyridin-3-yl)-1,3-thiazol-5-yl]phenyl}quinoline,
- 2-(3-{5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-2-yl}phenyl)propan-2-ol,
- 3-{5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-2-yl}benzoic acid,
- 2-{5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-2-yl}propan-2-ol,
- 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{2-(3-methyl-1,2,4-oxadiazol-5-yl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-5-yl}phenyl)quinoline,
- 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-5-yl}phenyl)quinoline,
- N*-cyclopropyl-5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazole-2-carboxamide,
- 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{2-(6-methyl-1-oxidopyridin-3-yl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-5-yl}phenyl)quinoline,

MC054PV


- 2-[4-(4-chlorophenyl)-5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-1,3-thiazol-2-yl]propan-2-ol,  
 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{2-methyl-4-[4-(methylsulfonyl)phenyl]-1,3-oxazol-5-yl}phenyl)quinoline,  
 5 2-{4-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-5-[4-(methylsulfonyl)phenyl]-1,3-thiazol-2-yl}propan-2-ol,  
 1,1,1-trifluoro-*N*-(5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-2-yl)methanesulfonamide,  
 2-[5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-4-pyridin-3-yl]-  
 10 1,3-thiazol-2-yl]propan-2-ol,  
 2-[5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-4-(1-oxidopyridin-3-yl)-1,3-thiazol-2-yl]propan-2-ol,  
 1-(4-chlorophenyl)-1-{4-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-5-[4-(methylsulfonyl)phenyl]-1,3-thiazol-2-yl}ethanol,  
 15 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{3-[4-(methylsulfonyl)phenyl]thien-2-yl}phenyl)quinoline,  
 1-(3''-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}-1,1':2',1''-terphenyl-4-yl)ethanone,  
 2-(3''-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}-1,1':2',1''-terphenyl-4-  
 20 yl]propan-2-ol,  
 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-{3-[5-(3-methyl-1,2,4-oxadiazol-5-yl)-1-oxidopyridin-3-yl]phenyl}quinoline,  
 5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-3-(3-methyl-1,2,4-oxadiazol-5-yl)pyridin-2(1*H*)-one,  
 25 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{6-[4-(methylsulfonyl)phenyl]imidazo[2,1-*b*][1,3]thiazol-5-yl}phenyl)quinoline,  
 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{2-[4-(methylsulfonyl)phenyl]imidazo[1,2-*a*]pyridin-3-yl}phenyl)quinoline  
 [3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}-4''-(methylthio)-1,1':2',1''-  
 30 terphenyl-4'-yl]methanol,  
 [3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}-4''-(methylsulfonyl)-1,1':2',1''-terphenyl-4'-yl]methanol,  
 2-[3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}-4''-(methylsulfonyl)-1,1':2',1''-terphenyl-4'-yl]propan-2-ol,

MC054PV

- 3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}-4''-(methylsulfonyl)-1,1':2',1''-terphenyl-4'-carboxylic acid,  
6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{2-[4-(methylsulfonyl)phenyl]-1-oxidopyridin-3-yl}phenyl)quinoline,  
5 8-{3-[3-(4-fluorophenyl)-1,2,4-oxadiazol-5-yl]phenyl}-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline,  
6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{4-[4-(methylthio)phenyl]pyridin-3-yl}phenyl)quinoline,  
3''-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}-1,1':2',1''-terphenyl-4-  
10 carboxylic acid,  
2-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-3-[4-(methylsulfonyl)phenyl]quinoxaline,  
6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{3-[4-(methylsulfonyl)phenyl]pyrazin-2-yl}phenyl)quinoline,  
15 5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-6-[4-(methylsulfonyl)phenyl]pyridin-2(1*H*)-one,  
1-methyl-5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-6-[4-(methylsulfonyl)phenyl]pyridin-2(1*H*)-one,  
8-(3-{6-methoxy-2-[4-(methylsulfonyl)phenyl]pyridin-3-yl}phenyl)-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline,  
20 8-(3-{6-(difluoromethoxy)-2-[4-(methylsulfonyl)phenyl]pyridin-3-yl}phenyl)-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline,  
8-(3-{6-[(4-fluorobenzyl)oxy]-2-[4-(methylsulfonyl)phenyl]pyridin-3-yl}phenyl)-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline,  
25 1-(4-fluorobenzyl)-5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-6-[4-(methylsulfonyl)phenyl]pyridin-2(1*H*)-one,  
5-(4-fluorophenyl)-6-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)pyridin-2(1*H*)-one,  
5-(4-fluorophenyl)-1-methyl-6-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)pyridin-2(1*H*)-one,  
30 8-{3-[3-(4-fluorophenyl)-6-methoxypyridin-2-yl]phenyl}-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline,  
and pharmaceutically acceptable salts thereof.

Further illustrating the invention are the compounds of the Formula:

MC054PV



wherein

R is selected from the group consisting of
Amino
2-biphenyl
3-biphenyl
N-(tert-butoxycarbonyl)-4-phenylpyrrolidin-3-yl
N-(tert-butoxycarbonyl)azetidin-3-yl
N-(tert-butoxycarbonyl)pyrrolidin-3-yl
3-chloro-4-fluorophenyl
4-chlorophenoxymethyl
2-chlorophenyl
4-chlorophenyl
Ethoxycarbonyl
furan-2-yl
furan-3-yl
imidazol-2-yl
indan-1-yl
indan-2-yl
1H-indol-2-yl
1H-indol-3-yl
1H-indol-4-yl
1H-indol-5-yl
1H-indol-6-yl
1H-indol-7-yl
Isoquinolin-1-yl

MC054PV

Isoquinolin-4-yl
Isoquinolin-5-yl
Isoquinolin-8-yl
isoxazol-3-yl
3-methoxycarbonylphenyl
4-methoxycarbonylphenyl
Methyl
1-methyl-1H-pyrazol-3-yl
1-methyl-1H-pyrazol-4-yl
1-methyl-1H-pyrazol-5-yl
2-methylphenyl
3-methylphenyl
4-methylphenyl
2-methylpyridin-5-yl
Methylsulfonylmethyl
2-methylsulfonylphenyl
3-methylsulfonylphenyl
4-methylsulfonylphenyl
morpholin-4-ylmethyl
Phenyl
Pyrazinyl
1H-pyrazol-3-yl
pyridin-2-yl
pyridin-3-yl
pyridin-4-yl
3-pyridinylmethyl
pyrimidin-2-yl
pyrimidin-4-yl
pyrimidin-5-yl
quinolin-4-yl
quinolin-5-yl
quinolin-8-yl
5,6,7,8-tetrahydro-1,8-naphthyridin-2-yl



6,7,8,9-tetrahydro-5H-benzo[a][7]annulen-5-yl
6,7,8,9-tetrahydro-5H-benzo[a][7]annulen-6-yl
Tetrahydrofuran-2-yl
1,2,3,4-tetrahydronaphthalen-1-yl
1,2,3,4-tetrahydronaphthalen-2-yl
1,3-thiazol-2-yl
1,3-thiazol-5-yl
thiophen-2-yl
thiophen-3-yl

And pharmaceutically acceptable salts thereof.

Further illustrating the invention are:

- 5 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{2-[3-(methylsulfonyl)phenyl]-4-phenyl-1,3-thiazol-5-yl}phenyl)quinoline,
- 2-[4-(3-chlorophenyl)-5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-1,3-thiazol-2-yl]propan-2-ol,
- 2-[4-(4-fluorophenyl)-5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-1,3-thiazol-2-yl]propan-2-ol,
- 10 8-{3-[4-(4-chlorophenyl)-2-quinolin-5-yl-1,3-thiazol-5-yl]phenyl}-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline,
- 2-{3-[4-(3-chlorophenyl)-5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-1,3-thiazol-2-yl]phenyl}propan-2-ol,
- 2-{3-[4-(3-chloro-4-fluorophenyl)-5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-1,3-thiazol-2-yl]phenyl}propan-2-ol,
- 15 2-{3-[4-[3,4-bis(difluoromethoxy)phenyl]-5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-1,3-thiazol-2-yl]phenyl}propan-2-ol,
- N*-(5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-2-yl)acetamide,
- 20 *N*-(5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-2-yl)pyridin-4-amine,
- 2-[5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-4-pyridin-4-yl-1,3-thiazol-2-yl]propan-2-ol,
- 2-[5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-4-(1-oxidopyridin-4-yl)-1,3-thiazol-2-yl]propan-2-ol,
- 25

MC054PV

- 2-[5-(4-chlorophenyl)-4-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-1,3-thiazol-2-yl]propan-2-ol,  
 2-{3-[4-(4-chlorophenyl)-5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-1,3-thiazol-2-yl]phenyl}propan-2-ol, and  
 5 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-{3-[5-(1*H*-tetraazol-5-yl)pyridin-3-yl]phenyl}quinoline,  
 and pharmaceutically acceptable salts thereof.

Further illustrating the invention are:

- 10 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{5-[4-(methylsulfonyl)phenyl]-1-oxidopyridin-3-yl}phenyl)quinoline,  
 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{5-[3-(methylsulfonyl)phenyl]-1-oxidopyridin-3-yl}phenyl)quinoline,  
 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{5-[2-(methylsulfonyl)phenyl]-1-oxidopyridin-3-yl}phenyl)quinoline,  
 15 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{2-[4-(methylsulfonyl)phenyl]pyridin-3-yl}phenyl)quinoline,  
 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-[3-(1-oxido-5-phenylpyridin-3-yl)phenyl]quinoline,  
 20 8-{3-[5-(3,5-dichlorophenyl)-1-oxidopyridin-3-yl]phenyl}-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline,  
 8-{3-[5-(3,4-dimethoxyphenyl)-1-oxidopyridin-3-yl]phenyl}-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline,  
 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-{3-[5-(5-methyl-1,2,4-oxadiazol-3-yl)-1-oxidopyridin-3-yl]phenyl}quinoline,  
 25 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-{3-[5-(5-methyl-1,3,4-oxadiazol-2-yl)-1-oxidopyridin-3-yl]phenyl}quinoline,  
 8-{3-[6-(benzyloxy)-5-(3-methyl-1,2,4-oxadiazol-5-yl)pyridin-3-yl]phenyl}-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline,  
 30 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{6-[4-(methylsulfonyl)phenyl]-1-oxidopyridin-3-yl}phenyl)quinoline,  
 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{5-[4-(methylsulfonyl)phenyl]-1-oxidopyridin-2-yl}phenyl)quinoline,  
 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-{3-[3-[4-(methylsulfonyl)phenyl]-5-(trifluoromethyl)pyridin-2-yl]phenyl}quinoline,  
 35

MC054PV

- 1-(4-chlorophenyl)-3-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)pyridin-2(1*H*)-one,  
*N*-isopropyl-5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-4-[4-(methylsulfonyl)phenyl]thiophene-2-carboxamide,  
 5 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(1,1':2',1''-terphenyl-3-yl)quinoline,  
 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-[4''-(methylthio)-1,1':2',1''-terphenyl-3-yl]quinoline,  
 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-[2'-(5-methyl-1,3,4-oxadiazol-2-yl)-1,1'-biphenyl-3-yl]quinoline,  
 10 methyl 3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}-4''-(methylsulfonyl)-1,1':2',1''-terphenyl-4'-carboxylate,  
 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(1,1':4',1''-terphenyl-3-yl)quinoline,  
 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(1,1':3',1''-terphenyl-3-yl)quinoline,  
 2-[5-(3'-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}-1,1'-biphenyl-2-yl)-1,3-thiazol-2-yl]propan-2-ol,  
 15 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-[3'-(1-oxidopyridin-4-yl)-1,1'-biphenyl-3-yl]quinoline,  
 and pharmaceutically acceptable salts thereof.

Further illustrating the invention are:

- 20 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{4-[4-(methylsulfonyl)phenyl]pyridin-3-yl}phenyl)quinoline,  
 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{4-[4-(methylsulfonyl)phenyl]-1-oxidopyridin-3-yl}phenyl)quinoline,  
 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{4-[3-(methylthio)phenyl]pyridin-3-yl}phenyl)quinoline,  
 25 8-[4',5'-difluoro-4''-(methylthio)-1,1':2',1''-terphenyl-3-yl]-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline,  
 8-[4',5'-difluoro-4''-(methylsulfonyl)-1,1':2',1''-terphenyl-3-yl]-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline,  
 30 8-(4''-fluoro-1,1':2',1''-terphenyl-3-yl)-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline,  
 6,7-dichloro-2-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-3-[4-(methylsulfonyl)phenyl]quinoxaline,  
 2-(4-chlorophenyl)-3-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)quinoxaline,

MC054PV

- 2-{4-[3-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)quinoxalin-2-yl]phenyl}propan-2-ol,  
 2-[3,4-bis(difluoromethoxy)phenyl]-3-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)quinoxaline,  
 5 4-[3-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)quinoxalin-2-yl]benzoic acid,  
*N*-cyclopropyl-4-[3-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)quinoxalin-2-yl]benzamide,  
 2-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-3-(4-methylphenyl)quinoxaline,  
 10 2-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-3-phenylquinoxaline,  
 2-(4-fluorophenyl)-3-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)quinoxaline,  
 15 2-{4-[3-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)pyrazin-2-yl]phenyl}propan-2-ol,  
 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-[3-[4-(methylthio)phenyl]pyrazin-2-yl]phenyl)quinoline,  
 8-[3-[3-(4-fluorophenyl)pyrazin-2-yl]phenyl]-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline,  
 20 8-(3-{2-(2-ethylpyridin-4-yl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-5-yl}phenyl)-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline,  
 2-(4-{5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-2-yl}phenyl)propan-2-ol,  
 25 and pharmaceutically acceptable salts thereof.

As used herein, "alkyl" as well as other groups having the prefix "alk" such as, for example, alkoxy, alkanoyl, alkenyl, alkynyl and the like, means carbon chains which may be linear or branched or combinations thereof. Examples of alkyl groups include methyl, ethyl, propyl, isopropyl, butyl, *sec*- and *tert*-butyl, pentyl, 30 hexyl, heptyl and the like. "Alkenyl", "alkynyl" and other like terms include carbon chains containing at least one unsaturated C-C bond.

The term "cycloalkyl" means carbocycles containing no heteroatoms, and includes mono-, bi- and tricyclic saturated carbocycles, as well as fused ring systems. Such fused ring systems can include one ring that is partially or fully

MC054PV

unsaturated such as a benzene ring to form fused ring systems such as benzofused carbocycles. Cycloalkyl includes such fused ring systems as spirofused ring systems. Examples of cycloalkyl include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, decahydronaphthalenyl, adamantanyl, indanyl, indenyl, fluorenyl, 1,2,3,4-  
5 tetrahydronaphthalenyl and the like. Similarly, "cycloalkenyl" means carbocycles containing no heteroatoms and at least one non-aromatic C-C double bond, and include mono-, bi- and tricyclic partially saturated carbocycles, as well as benzofused cycloalkenes. Examples of cycloalkenyl include cyclohexenyl, indenyl, and the like.

10 The term "cycloalkyloxy" unless specifically stated otherwise includes a cycloalkyl group connected to the oxy connecting atom.

The term "alkoxy" unless specifically stated otherwise includes an alkyl group connected to the oxy connecting atom.

The term "aryl" unless specifically stated otherwise includes multiple fused ring systems as well as single ring systems such as, for example, phenyl or  
15 naphthyl. In the case of single rings "aryl" shall include only aromatic carbocycles. In the case of multiple rings fused together, "aryl" shall include carbocycles where at least one of the rings is an aromatic carbocycle such as in indanyl, 1,2,3,4- tetrahydronaphthalene and 6,7,8,9-tertrhydro-5H-benzo[a]annulenyl.

20 The term "aryloxy" unless specifically stated otherwise includes multiple fused ring systems as mentioned above as well as single ring systems such as, for example, phenyl or naphthyl, connected through the oxy connecting atom to the connecting site.

The term "C<sub>0</sub>-C<sub>6</sub>alkyl" includes alkyls containing 6, 5, 4, 3, 2, 1, or no carbon atoms. An alkyl with no carbon atoms is a hydrogen atom substituent when  
25 the alkyl is a terminus moiety. An alkyl with no carbon atoms is a direct bond when the alkyl is a bridging moiety.

The term "heterocyclo" unless specifically stated otherwise includes one or more O, S, or N atoms. For example, heterocycles include ring systems that contain one or more O, S, or N atoms in the ring, including mixtures of such atoms.  
30 The heteroatoms replace ring carbon atoms. Thus, for example, a heterocyclo of 5 ring members is a five membered ring containing from zero to 4 carbon atoms. Moreover, for purposes of this specification, heterocycles includes rings substituted with "oxo", such as pyridin-2-one.

35 Examples of heterocycles include, for example, pyridinyl, pyridin-one-yl, quinolinyl, isoquinolinyl, pyridazinyl, pyrimidinyl, pyrazinyl, quinoxalinyl, furyl,

MC054PV

benzofuryl, dibenzofuryl, thienyl, benzothienyl, pyrrolyl, indolyl, pyrazolyl, indazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, imidazolyl, benzimidazolyl, oxadiazolyl, thiadiazolyl, triazolyl, tetrazolyl.

5 The term "heterocyclooxy" unless specifically stated otherwise describes a heterocyclo group connected through an oxy connecting atom to the connecting site.

Examples of heterocyclo(C<sub>1-6</sub>)alkyl include, for example, furylmethyl, furylethyl, thienylmethyl, thienylethyl, pyrazolylmethyl, oxazolylmethyl, oxazolylethyl, isoxazolylmethyl, thiazolylmethyl, thiazolylethyl, imidazolylmethyl, 10 imidazolylethyl, benzimidazolylmethyl, oxadiazolylmethyl, oxadiazolylethyl, thiadiazolylmethyl, thiadiazolylethyl, triazolylmethyl, triazolylethyl, tetrazolylmethyl, tetrazolylethyl, pyridinylmethyl, pyridinylethyl, pyridazinylmethyl, pyrimidinylmethyl, pyrazinylmethyl, quinolinylmethyl, isoquinolinylmethyl and quinoxalinylmethyl.

15 Other examples of heterocycloC<sub>3-7</sub>alkyl include, for example, azetidiny, pyrrolidinyl, piperidinyl, perhydroazepinyl, piperazinyl, morpholinyl, tetrahydrofuranyl, imidazolinyl, pyrrolidin-2-one, piperidin-2-one, and thiomorpholinyl.

The term "*N*-heterocycloC<sub>4-7</sub>alkyl" describes nonaryl heterocyclic compounds having 3-6 carbon atoms and one nitrogen atom forming the ring. 20 Examples include azetidiny, pyrrolidinyl, piperidinyl, and perhydroazepinyl.

Examples of aryl(C<sub>1-6</sub>)alkyl include, for example, phenyl(C<sub>1-6</sub>)alkyl, and naphthyl(C<sub>1-6</sub>)alkyl.

Examples of heterocycloC<sub>3-7</sub>alkylcarbonyl(C<sub>1-6</sub>)alkyl include, for 25 example, azetidiny carbonyl(C<sub>1-6</sub>)alkyl, pyrrolidinyl carbonyl(C<sub>1-6</sub>)alkyl, piperidinyl carbonyl(C<sub>1-6</sub>)alkyl, piperazinyl carbonyl(C<sub>1-6</sub>)alkyl, morpholinyl carbonyl(C<sub>1-6</sub>)alkyl, and thiomorpholinyl carbonyl(C<sub>1-6</sub>)alkyl.

The term "amine" unless specifically stated otherwise includes primary, secondary and tertiary amines.

30 Unless otherwise stated, the term "carbamoyl" is used to include -NHC(O)OC<sub>1-4</sub>alkyl, and -OC(O)NHC<sub>1-4</sub>alkyl.

The term "halogen" includes fluorine, chlorine, bromine and iodine atoms.

35 The term "optionally substituted" is intended to include both substituted and unsubstituted. Thus, for example, optionally substituted aryl could

MC054PV

represent a pentafluorophenyl or a phenyl ring. Further, the substitution can be made at any of the groups. For example, substituted aryl(C<sub>1-6</sub>)alkyl includes substitution on the aryl group as well as substitution on the alkyl group.

5 The term "oxide" of heteroaryl groups is used in the ordinary well-known chemical sense and include, for example, *N*-oxides of nitrogen heteroatoms.

Compounds described herein contain one or more double bonds and may thus give rise to *cis/trans* isomers as well as other conformational isomers. The present invention includes all such possible isomers as well as mixtures of such isomers.

10 Compounds described herein can contain one or more asymmetric centers and may thus give rise to diastereomers and optical isomers. The present invention includes all such possible diastereomers as well as their racemic mixtures, their substantially pure resolved enantiomers, all possible geometric isomers, and pharmaceutically acceptable salts thereof. The above Formula I is shown without a  
15 definitive stereochemistry at certain positions. The present invention includes all stereoisomers of Formula I and pharmaceutically acceptable salts thereof. Further, mixtures of stereoisomers as well as isolated specific stereoisomers are also included. During the course of the synthetic procedures used to prepare such compounds, or in using racemization or epimerization procedures known to those skilled in the art, the  
20 products of such procedures can be mixtures of stereoisomers.

The term "pharmaceutically acceptable salts" refers to salts prepared from pharmaceutically acceptable non-toxic bases or acids. When the compound of the present invention is acidic, its corresponding salt can be conveniently prepared from pharmaceutically acceptable non-toxic bases, including inorganic bases and  
25 organic bases. Salts derived from such inorganic bases include aluminum, ammonium, calcium, copper (ic and ous), ferric, ferrous, lithium, magnesium, manganese (ic and ous), potassium, sodium, zinc and the like salts. Particularly preferred are the ammonium, calcium, magnesium, potassium and sodium salts. Salts derived from pharmaceutically acceptable organic non-toxic bases include salts of  
30 primary, secondary, and tertiary amines, as well as cyclic amines and substituted amines such as naturally occurring and synthesized substituted amines. Other pharmaceutically acceptable organic non-toxic bases from which salts can be formed include ion exchange resins such as, for example, arginine, betaine, caffeine, choline, *N,N'*-dibenzylethylenediamine, diethylamine, 2-diethylaminoethanol, 2-  
35 dimethylaminoethanol, ethanolamine, ethylenediamine, *N*-ethylmorpholine, *N*-

MC054PV

ethylpiperidine, glucamine, glucosamine, histidine, hydrabamine, isopropylamine, lysine, methylglucamine, morpholine, piperazine, piperidine, polyamine resins, procaine, purines, theobromine, triethylamine, trimethylamine, tripropylamine, tromethamine and the like.

5           When the compound of the present invention is basic, its corresponding salt can be conveniently prepared from pharmaceutically acceptable non-toxic acids, including inorganic and organic acids. Such acids include, for example, acetic, benzenesulfonic, benzoic, camphorsulfonic, citric, ethanesulfonic, fumaric, gluconic, glutamic, hydrobromic, hydrochloric, isethionic, lactic, maleic, 10   malic, mandelic, methanesulfonic, mucic, nitric, pamoic, pantothenic, phosphoric, succinic, sulfuric, tartaric, p-toluenesulfonic acid and the like. Particularly preferred are benzenesulfonic, citric, hydrobromic, hydrochloric, maleic, phosphoric, sulfuric, and tartaric acids.

15           The pharmaceutical compositions of the present invention comprise a compound represented by Formula I (or pharmaceutically acceptable salts thereof) as an active ingredient, a pharmaceutically acceptable carrier and optionally other therapeutic ingredients or adjuvants. Such additional therapeutic ingredients include, for example, i) Leukotriene receptor antagonists, ii) Leukotriene biosynthesis inhibitors, iii) corticosteroids, iv) H1 receptor antagonists, v) beta 2 adrenoceptor 20   agonists, vi) COX-2 selective inhibitors, vii) statins, viii) non-steroidal anti-inflammatory drugs ("NSAID"), and ix) M2/M3 antagonists. The compositions include compositions suitable for oral, rectal, topical, and parenteral (including subcutaneous, intramuscular, and intravenous) administration, although the most suitable route in any given case will depend on the particular host, and nature and 25   severity of the conditions for which the active ingredient is being administered. The pharmaceutical compositions may be conveniently presented in unit dosage form and prepared by any of the methods well known in the art of pharmacy.

          Creams, ointments, jellies, solutions, or suspensions containing the compound of Formula I can be employed for topical use. Mouth washes and gargles 30   are included within the scope of topical use for the purposes of this invention.

          Dosage levels from about 0.01mg/kg to about 140mg/kg of body weight per day are useful in the treatment of conditions such as asthma, chronic bronchitis, chronic obstructive pulmonary disease (COPD), eosinophilic granuloma, psoriasis and other benign or malignant proliferative skin diseases, endotoxic shock 35   (and associated conditions such as laminitis and colic in horses), septic shock,



MC054PV

ulcerative colitis, Crohn's disease, reperfusion injury of the myocardium and brain, inflammatory arthritis, chronic glomerulonephritis, atopic dermatitis, urticaria, adult respiratory distress syndrome, chronic obstructive pulmonary disease in animals, diabetes insipidus, allergic rhinitis, allergic conjunctivitis, vernal conjunctivitis, arterial restenosis, orthrosclerosis, atherosclerosis, neurogenic inflammation, pain, cough, rheumatoid arthritis, ankylosing spondylitis, transplant rejection and graft versus host disease, hypersecretion of gastric acid, bacterial, fungal or viral induced sepsis or septic shock, inflammation and cytokine-mediated chronic tissue degeneration, osteoarthritis, cancer, cachexia, muscle wasting, depression, memory impairment, tumour growth and cancerous invasion of normal tissues which are responsive to PDE4 inhibition, or alternatively about 0.5mg to about 7g per patient per day. For example, inflammation may be effectively treated by the administration of from about 0.01mg to 50mg of the compound per kilogram of body weight per day, or alternatively about 0.5mg to about 3.5g per patient per day. Further, it is understood that the PDE4 inhibiting compounds of this invention can be administered at prophylactically effective dosage levels to prevent the above-recited conditions.

The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. For example, a formulation intended for the oral administration to humans may conveniently contain from about 0.5mg to about 5g of active agent, compounded with an appropriate and convenient amount of carrier material which may vary from about 5 to about 95 percent of the total composition. Unit dosage forms will generally contain between from about 1mg to about 500mg of the active ingredient, typically 25mg, 50mg, 100mg, 200mg, 300mg, 400mg, 500mg, 600mg, 800mg or 1000mg.

It is understood, however, that the specific dose level for any particular patient will depend upon a variety of factors including the age, body weight, general health, sex, diet, time of administration, route of administration, rate of excretion, drug combination and the severity of the particular disease undergoing therapy.

In practice, the compounds represented by Formula I, or pharmaceutically acceptable salts thereof, of this invention can be combined as the active ingredient in intimate admixture with a pharmaceutical carrier according to conventional pharmaceutical compounding techniques. The carrier may take a wide variety of forms depending on the form of preparation desired for administration, e.g., oral or parenteral (including intravenous). Thus, the pharmaceutical compositions of

MC054PV

the present invention can be presented as discrete units suitable for oral administration such as capsules, cachets or tablets each containing a predetermined amount of the active ingredient. Further, the compositions can be presented as a powder, as granules, as a solution, as a suspension in an aqueous liquid, as a non-aqueous liquid, as an oil-in-water emulsion or as a water-in-oil liquid emulsion. In addition to the common dosage forms set out above, the compound represented by Formula I, or pharmaceutically acceptable salts thereof, may also be administered by controlled release means and/or delivery devices. The compositions may be prepared by any of the methods of pharmacy. In general, such methods include a step of bringing into association the active ingredient with the carrier that constitutes one or more necessary ingredients. In general, the compositions are prepared by uniformly and intimately admixing the active ingredient with liquid carriers or finely divided solid carriers or both. The product can then be conveniently shaped into the desired presentation.

Thus, the pharmaceutical compositions of this invention may include a pharmaceutically acceptable carrier and a compound or a pharmaceutically acceptable salt of Formula I. The compounds of Formula I, or pharmaceutically acceptable salts thereof, can also be included in pharmaceutical compositions in combination with one or more other therapeutically active compounds.

The pharmaceutical carrier employed can be, for example, a solid, liquid, or gas. Examples of solid carriers include lactose, terra alba, sucrose, talc, gelatin, agar, pectin, acacia, magnesium stearate, and stearic acid. Examples of liquid carriers are sugar syrup, peanut oil, olive oil, and water. Examples of gaseous carriers include carbon dioxide and nitrogen.

In preparing the compositions for oral dosage form, any convenient pharmaceutical media may be employed. For example, water, glycols, oils, alcohols, flavoring agents, preservatives, coloring agents and the like may be used to form oral liquid preparations such as suspensions, elixirs and solutions; while carriers such as starches, sugars, microcrystalline cellulose, diluents, granulating agents, lubricants, binders, disintegrating agents, and the like may be used to form oral solid preparations such as powders, capsules and tablets. Because of their ease of administration, tablets and capsules are the preferred oral dosage units whereby solid pharmaceutical carriers are employed. Optionally, tablets may be coated by standard aqueous or nonaqueous techniques

A tablet containing the composition of this invention may be prepared by compression or molding, optionally with one or more accessory ingredients or adjuvants. Compressed tablets may be prepared by compressing, in a suitable machine, the active ingredient in a free-flowing form such as powder or granules, optionally mixed with a binder, lubricant, inert diluent, surface active or dispersing agent. Molded tablets may be made by molding in a suitable machine, a mixture of the powdered compound moistened with an inert liquid diluent. Each tablet preferably contains from about 0.1mg to about 500mg of the active ingredient and each cachet or capsule preferably containing from about 0.1mg to about 500mg of the active ingredient.

Pharmaceutical compositions of the present invention suitable for parenteral administration may be prepared as solutions or suspensions of the active compounds in water. A suitable surfactant can be included such as, for example, hydroxypropylcellulose. Dispersions can also be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof in oils. Further, a preservative can be included to prevent the detrimental growth of microorganisms.

Pharmaceutical compositions of the present invention suitable for injectable use include sterile aqueous solutions or dispersions. Furthermore, the compositions can be in the form of sterile powders for the extemporaneous preparation of such sterile injectable solutions or dispersions. In all cases, the final injectable form must be sterile and must be effectively fluid for easy syringability. The pharmaceutical compositions must be stable under the conditions of manufacture and storage; thus, preferably should be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (e.g. glycerol, propylene glycol and liquid polyethylene glycol), vegetable oils, and suitable mixtures thereof.

Pharmaceutical compositions of the present invention can be in a form suitable for topical use such as, for example, an aerosol, cream, ointment, lotion, dusting powder, or the like. Further, the compositions can be in a form suitable for use in transdermal devices. These formulations may be prepared, utilizing a compound represented by Formula I of this invention, or pharmaceutically acceptable salts thereof, via conventional processing methods. As an example, a cream or ointment is prepared by mixing hydrophilic material and water, together with about 5 wt% to about 10 wt% of the compound, to produce a cream or ointment having a desired consistency.

MC054PY

Pharmaceutical compositions of this invention can be in a form suitable for rectal administration wherein the carrier is a solid. It is preferable that the mixture forms unit dose suppositories. Suitable carriers include cocoa butter and other materials commonly used in the art. The suppositories may be conveniently  
5 formed by first admixing the composition with the softened or melted carrier(s) followed by chilling and shaping in moulds.

In addition to the aforementioned carrier ingredients, the pharmaceutical formulations described above may include, as appropriate, one or more additional carrier ingredients such as diluents, buffers, flavoring agents, binders,  
10 surface-active agents, thickeners, lubricants, preservatives (including anti-oxidants) and the like. Furthermore, other adjuvants can be included to render the formulation isotonic with the blood of the intended recipient. Compositions containing a compound described by Formula I, or pharmaceutically acceptable salts thereof, may also be prepared in powder or liquid concentrate form.

15 The compounds and pharmaceutical compositions of this invention have been found to exhibit biological activity as PDE4 inhibitors. Accordingly, another aspect of the invention is the treatment in mammals of, for example, asthma, chronic bronchitis, chronic obstructive pulmonary disease (COPD), eosinophilic granuloma, psoriasis and other benign or malignant proliferative skin diseases,  
20 endotoxic shock (and associated conditions such as laminitis and colic in horses), septic shock, celiac sprue, ulcerative colitis, Crohn's disease, reperfusion injury of the myocardium and brain, inflammatory arthritis, chronic glomerulonephritis, atopic dermatitis, urticaria, adult respiratory distress syndrome, chronic obstructive pulmonary disease in animals, diabetes insipidus, allergic rhinitis, allergic  
25 conjunctivitis, vernal conjunctivitis, arterial restenosis, orthosclerosis, atherosclerosis, neurogenic inflammation, pain, cough, rheumatoid arthritis, ankylosing spondylitis, transplant rejection and graft versus host disease, hypersecretion of gastric acid, bacterial, fungal or viral induced sepsis or septic shock, inflammation and cytokine-mediated chronic tissue degeneration, osteoarthritis,  
30 cancer, cachexia, muscle wasting, depression, memory impairment, tumour growth and cancerous invasion of normal tissues – maladies that are amenable to amelioration through inhibition of the PDE4 isoenzyme and the resulting elevated cAMP levels – by the administration of an effective amount of the compounds of this invention.

35 In addition to being useful in the treatment of cognitive deficit (such as memory impairment, mentioned elsewhere in this specification) due to psychological

MC054PV

dysfunction, neurological deficit (such as stroke) or psychiatric dysfunction, the compounds of the invention are useful for enhancing cognition in a healthy subject (enhancing memory, learning, retention, recall, awareness and judgement).

5 The term "mammals" includes humans, as well as other animals such as; for example, dogs, cats, horses, pigs, and cattle. Accordingly, it is understood that the treatment of mammals other than humans is the treatment of clinical correlating afflictions to those above recited examples that are human afflictions.

Further, as described above, the compound of this invention can be utilized in combination with other therapeutic compounds. In particular, the combinations of the PDE4 inhibiting compound of this invention can be advantageously used in combination with i) Leukotriene receptor antagonists, ii) Leukotriene biosynthesis inhibitors, iii) corticosteroids, iv) H1 receptor antagonists, v) beta 2 adrenoceptor agonists, vi) COX-2 selective inhibitors, vii) statins, viii) non-steroidal anti-inflammatory drugs ("NSAID"), and ix) M2/M3 antagonists.

15 The abbreviations used herein have the following tabulated meanings. Abbreviations not tabulated below have their meanings as commonly used unless specifically stated otherwise.

Ac	=	acetyl
Bn	=	benzyl
CAMP		cyclic adenosine-3',5'-monophosphate
DBU	=	1,8-diazabicyclo[5.4.0]undec-7-ene
DIBAL	=	diisobutylaluminum hydride
DMAP	=	4-(dimethylamino)pyridine
DMF	=	N,N-dimethylformamide
Et <sub>3</sub> N	=	triethylamine
GST		glutathione transferase
HMDS		hexamethyldisilazide
LDA	=	lithium diisopropylamide
m-CPBA	=	metachloroperbenzoic acid
MMPP	=	monoperoxyphthalic acid
MPPM	=	monoperoxyphthalic acid, magnesium salt 6H <sub>2</sub> O
Ms	=	methanesulfonyl = mesyl = SO <sub>2</sub> Me
MsO	=	methanesulfonate = mesylate

NSAID =	non-steroidal anti-inflammatory drug
o-Tol =	ortho-tolyl
OXONE® =	2KHSO <sub>5</sub> •KHSO <sub>4</sub> •K <sub>2</sub> SO <sub>4</sub>
PCC =	pyridinium chlorochromate
PDC =	pyridinium dichromate
PDE	phosphodiesterase
Ph =	phenyl
Phe =	benzenediyl
PMB =	para-methoxybenzyl
Pye =	pyridinediyl
r.t. =	room temperature
Rac. =	racemic
SAM =	aminosulfonyl or sulfonamide or SO <sub>2</sub> NH <sub>2</sub>
SEM =	2-(trimethylsilyl)ethoxymethoxy
SPA =	scintillation proximity assay
TBAF =	tetra-n-butylammonium fluoride
Th =	2- or 3-thienyl
TFA =	trifluoroacetic acid
TFAA =	trifluoroacetic acid anhydride
THF =	tetrahydrofuran
Thi =	thiophenediyl
TLC =	thin layer chromatography
TMS-CN =	trimethylsilyl cyanide
TMSI	trimethylsilyl iodide
Tz =	1H (or 2H)-tetrazol-5-yl
C <sub>3</sub> H <sub>5</sub> =	allyl

#### ALKYL GROUP ABBREVIATIONS

Me =	Methyl
Et =	Ethyl
<i>n</i> -Pr =	Normal propyl
<i>i</i> -Pr =	Isopropyl

<i>n</i> -Bu	=	Normal butyl
<i>i</i> -Bu	=	Isobutyl
<i>s</i> -Bu	=	Secondary butyl
<i>t</i> -Bu	=	Tertiary butyl
c-Pr	=	Cyclopropyl
c-Bu	=	Cyclobutyl
c-Pen	=	Cyclopentyl
c-Hex	=	Cyclohexyl

### ASSAYS DEMONSTRATING BIOLOGICAL ACTIVITY

#### LPS AND FMLP-INDUCED TNF- $\alpha$ AND LTB<sub>4</sub> ASSAYS IN HUMAN WHOLE BLOOD

Whole blood provides a protein and cell-rich milieu appropriate for the study of biochemical efficacy of anti-inflammatory compounds such as PDE4-selective inhibitors. Normal non-stimulated human blood does not contain detectable levels of TNF- $\alpha$  and LTB<sub>4</sub>. Upon stimulation with LPS, activated monocytes express and secrete TNF- $\alpha$  up to 8 hours and plasma levels remain stable for 24 hours. Published studies have shown that inhibition of TNF- $\alpha$  by increasing intracellular cAMP via PDE4 inhibition and/or enhanced adenylyl cyclase activity occurs at the transcriptional level. LTB<sub>4</sub> synthesis is also sensitive to levels of intracellular cAMP and can be completely inhibited by PDE4-selective inhibitors. As there is little LTB<sub>4</sub> produced during a 24 hour LPS stimulation of whole blood, an additional LPS stimulation followed by fMLP challenge of human whole blood is necessary for LTB<sub>4</sub> synthesis by activated neutrophils. Thus, by using the same blood sample, it is possible to evaluate the potency of a compound on two surrogate markers of PDE4 activity in the whole blood by the following procedure.

Fresh blood was collected in heparinized tubes by venipuncture from healthy human volunteers (male and female). These subjects had no apparent inflammatory conditions and had not taken any NSAIDs for at least 4 days prior to blood collection. 500 $\mu$ L aliquots of blood were pre-incubated with either 2 $\mu$ L of vehicle (DMSO) or 2 $\mu$ L of test compound at varying concentrations for 15 minutes at 37°C. This was followed by the addition of either 10 $\mu$ L vehicle (PBS) as blanks or 10 $\mu$ L LPS (1 $\mu$ g/mL final concentration, #L-2630 (Sigma Chemical Co., St. Louis,

MC054PV

- MO) from *E. coli*, serotype 0111:B4; diluted in 0.1% w/v BSA (in PBS)). After 24 hours of incubation at 37°C, another 10µL of PBS (blank) or 10µL of LPS (1µg/mL final concentration) was added to blood and incubated for 30 minutes at 37°C. The blood was then challenged with either 10µL of PBS (blank) or 10µL of fMLP (1µM final concentration, #F-3506 (Sigma); diluted in 1% w/v BSA (in PBS)) for 15 minutes at 37°C. The blood samples were centrifuged at 1500xg for 10 minutes at 4°C to obtain plasma. A 50µL aliquot of plasma was mixed with 200µL methanol for protein precipitation and centrifuged as above. The supernatant was assayed for LTB<sub>4</sub> using an enzyme immunoassay kit (#520111 from Cayman Chemical Co., Ann Arbor, MI) according to the manufacturer's procedure. TNF-α was assayed in diluted plasma (in PBS) using an ELISA kit (Cistron Biotechnology, Pine Brook, NJ) according to manufacturer's procedure. The IC<sub>50</sub> values of selected examples:

Example:	IC <sub>50</sub> (µM)
115	0.61
167	0.139
1	0.155
19	0.046
24	0.054
41	0.044
69	0.025
78	0.12
82	0.156
88	0.144
94	1.04
100	0.220
157	0.136

15

#### ANTI-ALLERGIC ACTIVITY *IN VIVO*

- Compounds of the invention have been tested for effects on an IgE-mediated allergic pulmonary inflammation induced by inhalation of antigen by sensitized guinea pigs. Guinea pigs were initially sensitized to ovalbumin under mild cyclophosphamide-induced immunosuppression, by intraperitoneal injection of

20



antigen in combinations with aluminum hydroxide and pertussis vaccine. Booster doses of antigen were given two and four weeks later. At six weeks, animals were challenged with aerosolized ovalbumin while under cover of an intraperitoneally administered anti-histamine agent (mepyramine). After a further 48h, bronchial alveolar lavages (BAL) were performed and the numbers of eosinophils and other leukocytes in the BAL fluids were counted. The lungs were also removed for histological examination for inflammatory damage. Administration of compounds of the Examples (0.001-10mg/kg i.p. or p.o.), up to three times during the 48h following antigen challenge, lead to a significant reduction in the eosinophilia and the accumulation of other inflammatory leukocytes. There was also less inflammatory damage in the lungs of animals treated with compounds of the Examples.

#### SPA BASED PDE ACTIVITY ASSAY PROTOCOL

Compounds which inhibit the hydrolysis of cAMP to AMP by the type-IV cAMP-specific phosphodiesterases were screened in a 96-well plate format as follows:

In a 96 well-plate at 30°C was added the test compound (dissolved in 2μL DMSO), 188μL of substrate buffer containing [2,8-<sup>3</sup>H] adenosine 3',5'-cyclic phosphate (cAMP, 100nM to 50μM), 10mM MgCl<sub>2</sub>, 1mM EDTA, 50mM Tris, pH 7.5. The reaction was initiated by the addition of 10μL of human recombinant PDE4 (the amount was controlled so that ~10% product was formed in 10min.). The reaction was stopped after 10min. by the addition of 1mg of PDE-SPA beads (Amersham Pharmacia Biotech, Inc., Piscataway, NJ). The product AMP generated was quantified on a Wallac Microbeta® 96-well plate counter (EG&G Wallac Co., Gaithersburg, MD). The signal in the absence of enzyme was defined as the background. 100% activity was defined as the signal detected in the presence of enzyme and DMSO with the background subtracted. Percentage of inhibition was calculated accordingly. IC<sub>50</sub> value was approximated with a non-linear regression fit using the standard 4-parameter/multiple binding sites equation from a ten point titration.

The IC<sub>50</sub> values of Examples in table below were determined with 100nM cAMP using the purified GST fusion protein of the human recombinant phosphodiesterase IVa (met-248) produced from a baculovirus/Sf-9 expression system. The IC<sub>50</sub> values of selected examples:

MC054PV

Example:	IC50 (nM)
115	2.2
167	0.4
1	0.3
19	1.0
24	0.4
41	1.0
69	0.6
78	0.4
82	3.1
88	1.6
94	4.3
100	4.1
157	1.7

The examples that follow are intended as an illustration of certain preferred embodiments of the invention and no limitation of the invention is implied.

5 Unless specifically stated otherwise, the experimental procedures were performed under the following conditions. All operations were carried out at room or ambient temperature - that is, at a temperature in the range of 18-25°C. Evaporation of solvent was carried out using a rotary evaporator under reduced pressure (600-4000pascals: 4.5-30mm. Hg) with a bath temperature of up to 60°C. The course of  
10 reactions was followed by thin layer chromatography (TLC) and reaction times are given for illustration only. Melting points are uncorrected and 'd' indicates decomposition. The melting points given are those obtained for the materials prepared as described. Polymorphism may result in isolation of materials with different melting points in some preparations. The structure and purity of all final  
15 products were assured by at least one of the following techniques: TLC, mass spectrometry, nuclear magnetic resonance (NMR) spectrometry or microanalytical data. Yields are given for illustration only. When given, NMR data is in the form of delta ( $\delta$ ) values for major diagnostic protons, given in parts per million (ppm) relative to tetramethylsilane (TMS) as internal standard, determined at 300MHz, 400MHz or  
20 500MHz using the indicated solvent. Conventional abbreviations used for signal

MC054PV

shape are: s. singlet; d. doublet; t. triplet; m. multiplet; br. broad; etc. In addition, "Ar" signifies an aromatic signal. Chemical symbols have their usual meanings; the following abbreviations have also been used: v (volume), w (weight), b.p. (boiling point), m.p. (melting point), L (liter(s)), mL (milliliters), g (gram(s)), mg (milligrams(s)), mol (moles), mmol (millimoles), eq (equivalent(s)).

### Methods of Synthesis

Compounds of the present invention can be prepared according to the following methods. The substituents are the same as in Formula I except where defined otherwise.

### METHODS OF SYNTHESIS


Compounds of the present invention can be prepared according to the following general methods. Reactions are typically run under nitrogen atmosphere at ambient temperature if not otherwise mention. Anhydrous solvent such as THF, DMF, Et<sub>2</sub>O, DME and Toluene are commercial grade. Reagents are commercial grade and were used without any purification. Flash chromatography is run on silica gel (230-400 mesh).

All 8-aryl-quinoline of the type I can be prepared (Scheme 1 and 2) using a Suzuki type coupling to build the biaryl moieties. In a typical Suzuki coupling reaction, all reagents except for the palladium catalyst are mixed in the appropriate solvent. The mixture is then degassed by refluxing for 15 min under nitrogen atmosphere then cooling to ambient temperature or by applying two to three vacuum / nitrogen sequence. The palladium catalyst is then added and the reaction mixture is stirred at the appropriate temperature until completion as monitored by TLC.

The substituents are the same as in Formula I except where defined otherwise. Compounds of the type I (Scheme 1) can be prepared in a two step one-pot manner by generating *in-situ* the boronate analog of 8-bromo quinoline II followed by a palladium catalyzed coupling with the appropriate biaryl III

MC054PV


Scheme1



- In most cases, compounds were prepared by the two procedures described in Scheme
- 5 2. A Suzuki type coupling between the 8-bromo-quinoline **II** and the bromo-phenyl-  
boronic acid **IV** produced to the common intermediate **V**. The latter can be coupled  
with either an aryl-stannane of type **VI** or a boronic acid of type **VII** to generate the  
desired compound **I**. Alternatively, the arylbromide **V** can be converted to the  
corresponding pinacole boronate **VIII** by a  $\text{PdCl}_2(\text{dppf})_2$  catalyzed coupling reaction.
- 10 with pinacole diborane. Subsequently, a Suzuki coupling of the boronate **VIII** with  
the appropriate heteroaryl halide or O-triflate **IX** will generate the desired compound  
**I**.


MC054PV

Scheme 2




- 5 Boronic acid of the type **XII** can be prepared (Scheme 3) by lithium-halogen exchange at low temperature in THF or Et<sub>2</sub>O on the corresponding heteroaryl bromide **X** followed by the addition of a trialkyl-boronate (B(OR)<sub>3</sub>). Hydrolysis, under acidic condition, of the resulting heteroaryl-boronate will generate the desired boronic acid **XII** (R=H). Likewise, lithium-halogen exchange or deprotonation at low temperature
- 10 in THF or Et<sub>2</sub>O followed by the addition of a trialkylstannyl-chloride (R<sub>3</sub>SnCl) generates the stannane of type **XI**. These can be used as precursors for biaryl of type **IX**.

Scheme 3




- The heterocyclic  $\text{Ar}_1$  groups can be constructed from the common aldehyde **XIV** or **Q-CHO** (scheme 4) prepared by Suzuki type coupling. The aldehyde **XIV** can lead to ethanone **XV** or ethanone **XVI** (scheme 5) by using a methodology described by M. Journet *et al.* in *Tetrahedron Lett.* 39, 1717 (1998). They can serve as precursor for pyridinone **XVII** and **XVIII** using ammonia and methyl propiolate as described for example in *J. Heterocyclic Chem.* 30, 1129 (1993). The ethanone **XV** or ethanone **XVI** can also be lead to alpha bromo analogs **XIX** and **XX** which are used for the formation of thiazole, oxazole **XXI**, **XXII** and other heterocycle like quinoxaline **XXIII**, imidazothiazole **XXIV** or imidazopyridinyl **XXV**.

Scheme 4



Scheme 5




5

8-Bromoquinoline II with various substituent at position 6 can be prepared by the Skraup synthesis (Org. React. 7, 59-98, 1953) using an appropriate aniline, an oxidizing agent such as 3-nitrophenylsulfonic acid and glycerol in strong acidic media such as sulfuric acid or methanesulfonic acid (Scheme 6).

10

MC054PV


## Scheme 6



5

The common quinoline intermediates were prepared using the following procedures.

## Quinoline\_1




10

## 8-Bromo-6-(1-methanesulfonyl-1-methyl-ethyl)-quinoline

Preparation of 8-Bromo-6-(1-methanesulfonyl-1-methyl-ethyl)-quinoline is described in Patent Publication US 6 410 563 B1 (2002).

15

## Quinoline\_2




## 3-{3-[1-methyl-1-(methanesulfonyl)ethyl]-1-naphthyl}benzaldehyde



MC054PV

Preparation of 3-{3-[1-methyl-1-(methanesulfonyl)ethyl]-1-naphthyl}benzaldehyde is described in Patent Publication US 6 410 563 B1 (2002).


### Quinoline\_3



### 8-(3-Bromo-phenyl)-6-(1-methanesulfonyl-1-methyl-ethyl)-quinoline

A mixture of Quinoline 2 (1.0 eq), 3-bromophenylboronic acid (1.05 eq),  $\text{Na}_2\text{CO}_3$  (2 M in  $\text{H}_2\text{O}$ ; 3.6 eq) and  $\text{Pd}(\text{PPh}_3)_4$  (0.03 eq) in DME (0.2 M) was stirred at  $80^\circ\text{C}$  for 8 h. The resulting mixture was cooled to room temperature and diluted with water under vigorous stirring. The resulting precipitate was filtered and dried. Flash chromatography (Hex:EtOAc; 2:3) and stirring in a mixture of  $\text{Et}_2\text{O}$  and  $\text{CH}_2\text{Cl}_2$  (10:1) yielded the title compound as a light yellow solid after filtration.

### Quinoline\_4




### 6-(1-Methanesulfonyl-1-methyl-ethyl)-8-[3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-quinoline

A mixture of Quinoline\_3 (1.0 eq), pinacole diborane ester (1.4 eq) and KOAc (3.5 eq) and  $\text{PdCl}_2(\text{dppf})_2$  (0.03 eq) in DMF (0.14 M) was stirred at  $60^\circ\text{C}$  for 24 h. An extra amount of pinacole diborane (0.3 eq), KOAc (1.05 eq) and  $\text{PdCl}_2(\text{dppf})_2$  (0.01 eq) were added and the mixture was stirred at  $60^\circ\text{C}$  for 24 h. The resulting mixture

MC054PV

was cooled to room temperature, diluted with EtOAc:Et<sub>2</sub>O (1:1). The organic phase was washed with water (3x), brine, dried over MgSO<sub>4</sub>, filtered and concentrated. Flash chromatography (CH<sub>2</sub>Cl<sub>2</sub>:EtOAc ; 9:1) and stirring in Et<sub>2</sub>O:EtOAc (10:1) afforded the title compound as a white solid. Subsequent flash chromatography (CH<sub>2</sub>Cl<sub>2</sub>:EtOAc ; 9:1) on the mother liquor and stirring in Et<sub>2</sub>O:EtOAc (10:1) afforded a second crop of the title compound containing 5 % of starting material.

**Quinoline\_5**

8-bromo-N-cyclopropylquinoline-6-carboxamide

**Step 1: 8-bromoquinoline-6-carboxylic acid**

4-Amino-3-bromo-benzoic acid (1 eq) and glycerol (2.4 eq) was added to sulfuric acid (6.4M, reaction conc. 0.6M) at 0°C. After 15 min, the bath was removed and 3-nitro-phenylsulfonic acid sodium salt (1.3 eq) was added portionwise. The reaction mixture was heated gradually until exothermic reaction started (between 110-140°C), the internal temperature was maintained below 170°C for 1h. The reaction mixture was cooled to 50°C and poured slowly into NH<sub>4</sub>OH-ice (2:1). The title product isolated by filtration, dissolved in warm 5% NH<sub>4</sub>OH/MeOH and diluted with CH<sub>2</sub>Cl<sub>2</sub>. This solution was filtered through silica gel eluting with (NH<sub>4</sub>OH, MeOH, CH<sub>2</sub>Cl<sub>2</sub>, 3:30:70). The residue was stirred in EtOH and the title product isolated by filtration.

**Step 2: 8-bromo-N-cyclopropylquinoline-6-carboxamide**


A solution of HATU (1.2 eq), 8-bromoquinoline-6-carboxylic acid (1 eq), diisopropylethylamine (2.5 eq) and cyclopropylamine (1.1 eq) was stirred at rt for 1h in DMF (0.2M). The reaction mixture was diluted with water, stirred for

MC054PV

2h and extracted with EtOAc. The organic extracts were washed with brine, dried over  $\text{MgSO}_4$ , filtered and concentrated. The residue was crystallized in EtOAc and title product isolated by filtration as a white powder.

5

#### Quinoline\_6



#### 2-(8-bromoquinolin-6-yl)-2-methylpropanenitrile

Preparation of 2-(8-bromoquinolin-6-yl)-2-methylpropanenitrile is described in Patent Publication US 6 410 563 B1 (2002).

#### General Method of Synthesis:

##### 15 Coupling-1; General Procedure for Aryl-Aryl coupling

A mixture of boronic acid or boronate ester (1.0 eq), aryl bromide (chloride or iodide) (1.1-2.0 eq.),  $\text{Na}_2\text{CO}_3$  (3.0-3.5 eq.; 2 M in  $\text{H}_2\text{O}$ ) and  $\text{PdCl}_2(\text{dppf})_2$  or  $\text{Pd}(\text{OAc})_2(\text{Ph}_3\text{P})_2$  or  $\text{Pd}(\text{Ph}_3\text{P})_4$  or  $\text{Pd}(1,3\text{-Bis}(2,4,6\text{-trimethylphenyl})\text{imidazolium chloride})_2$  (0.05 eq.) in DME (or n-Propanol or Dioxane or DMF) (0.05-0.2 M) was stirred at 60-80°C for 1-18 h. The mixture was cooled to room temperature and concentrated. Flash chromatography afforded the title compound. (Coupling using Aryl Iodide generally proceeds at lower temperature 20-60°C).

##### 25 Coupling-2; General Procedure for Aryl-Aryl coupling

A mixture of boronic acid or boronate ester (1.0 eq), aryl bromide (or chloride or iodide) (1.1-2.0 eq),  $\text{CsF}$  (2.5 eq) or aqueous  $\text{Na}_2\text{CO}_3$  (2M),  $\text{Pd}_2(\text{dba})_3$  (1.5 %) and  $(t\text{-Bu})_3\text{P}$  (4.5%) in THF or dioxane (0.05-0.2 M) was stirred at 20°C for 12-18 h. The mixture was concentrated, diluted with EtOAc and water then extracted. Flash chromatography afforded the title compound.

30

##### Coupling-3; General Procedure for Aryl-Aryl coupling

MC054PV

A mixture of Aryltributyltin (1-2 eq), aryl bromide (1-2 eq), CuI (0.05 eq), PdCl<sub>2</sub>(dppf) (0.05 eq) in dioxane (0.05-0.2 M) was stirred at 100°C for 12-18 h. The mixture was concentrated, diluted with EtOAc and water then extracted. Flash chromatography afforded the title compound.

5

**Oxid-1; General Procedure for oxidation of sulfide to sulfone**

To a solution Sulfide (1.0 eq.) in THF:MeOH:H<sub>2</sub>O (2:2:1) (or alternatively replacing water with saturated aqueous NaHCO<sub>3</sub>), was added Oxone (2.2 eq.). The mixture was stirred for 12 h at room temperature, poured in water and extracted with EtOAc (2x).

10 The combined organic extracts were washed with aqueous NaHCO<sub>3</sub>, brine, dried over MgSO<sub>4</sub> or Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated. Flash chromatography afforded the title compound.

**Oxid-2: General Procedure for oxidation of Pyridinyl to Pyridinyl-N-oxide:**

15 To a solution of Pyridinyl (1.0 eq.) in CH<sub>2</sub>Cl<sub>2</sub> (0.1 M) was added *m*-CPBA (1.3 eq.). The mixture was stirred at room temperature for 12 h, quenched with Ca(OH)<sub>2</sub> (0.7eq) diluted with CH<sub>2</sub>Cl<sub>2</sub> and filtered. The filtrate was concentrated and flash chromatography afforded the title compound.

Alternatively, the reaction mixture was quenched with EtOAc and aqueous NaHCO<sub>3</sub>, the combined organic extracts were washed with aqueous NaHCO<sub>3</sub>, brine, dried over MgSO<sub>4</sub> or Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated. Flash chromatography afforded the title compound.

20

**Cer-1; General Procedure for Grignard type addition on ester and ketone using CeCl<sub>3</sub>**

25

A heterogeneous solution of CeCl<sub>3</sub> (1.5 eq) and ester (1 eq) in THF (0.1M) is sonicated at room temperature for 15 minutes. The resulting suspension is cooled to -78°C and RMgX (Cl, Br, I, 6 eq) is added. The mixture is let to warm to -25°C over a 2 hours period. The reaction is quenched with saturated solution of NH<sub>4</sub>Cl and diluted with ethyl acetate. The organic phase is extracted, washed with brine and dry over MgSO<sub>4</sub> and concentrated. The residu was purified by flash chromatography (Hexane:EtOAc) or precipitated with ether/ethyl acetate to afforded the title compound.

30

MC054PV

Alternatively ketones can be used as starting material with 3 eq of Grignard reagent (RMgX).

**Thio-1: General Procedure for conversion of nitrile to thioamide:**


- 5 Heating nitrile derivative in MeOH at reflux with ammonium sulfide as described in SPYCHALA, J.; Synth Commun. 1997, 27 (19), 3431-3440. The product is generally purified by filtration, wash with water and air dried.

Bi-Aryl intermediates were prepared using the following procedures.

10

**Pyr\_1**

**3-bromo-5-[4-(methylsulfonyl)phenyl]pyridine 1-oxide**



- 15 **Step 1 : 3-bromo-5-[4-(methylthio)phenyl]pyridine**

Prepared according to the procedure **Coupling-2** using [4-(methylthio)phenyl]boronic acid and 3,5-dibromopyridine as starting material. Flash chromatography (Hex:EtOAc ; 95:5-85:15) afforded the title compound


20

**Step 2 : 3-bromo-5-[4-(methylsulfonyl)phenyl]pyridine 1-oxide**

Prepared according to the procedure **Oxid-1** and using oxone in excess (5 eq). The residue was stirred vigorously in EtOAc then isolated by filtration.

25

MC054PV

**Pyr\_2****3-bromo-5-[3-(methylsulfonyl)phenyl]pyridine 1-oxide****5 Step 1 : 3-bromo-5-[3-(methylthio)phenyl]pyridine**


Prepared according to the procedure **Coupling-2** using [3-(methylthio)phenyl]boronic acid and 3,5-dibromopyridine as starting material. Flash chromatography (Hex:EtOAc ; 95:5-85:15) afforded the title compound.

10

**Step 2 : 3-bromo-5-[3-(methylsulfonyl)phenyl]pyridine 1-oxide**

Prepared according to the procedure **Oxid-2** and using mCPBA (3.2 eq). The residue was stirred vigorously in Et<sub>2</sub>O then isolated by filtration.

15


**Pyr\_3****3-bromo-5-[2-(methylsulfonyl)phenyl]pyridine 1-oxide**

20 Prepared according to the procedure described for **Pyr\_2** using [2-(methylthio)phenyl]boronic acid as starting material. The residue was stirred vigorously in Et<sub>2</sub>O then isolated by filtration.

MC054PV

## Pyr\_4

## 3-iodo-2-[4-(methylsulfonyl)phenyl]pyridine



## 5 Step 1 : 3-amino-2-[4-(methylthio)phenyl]pyridine

Prepared according to the procedure **Coupling-1** using [4-(methylthio)phenyl]boronic acid and 2-amino-1-chloropyridine as starting material. Flash chromatography (Hex:EtOAc ; 60:40-50:50) afforded the title compound.

10

## Step 2 : 3-amino-2- [4-(methylsulfonyl)phenyl]pyridine

Prepared according to the procedure **Oxid-1**. Flash chromatography (Hex:EtOAc ; 50:50-20:80) afforded the title compound as a white solid.


15

## Step 3 : 3-Iodo-2- [4-(methylsulfonyl)phenyl]pyridine


To a solution of 3-amino-2- [4-(methylsulfonyl)phenyl]pyridine (1.0 eq.) in AcOH:CH<sub>2</sub>Cl<sub>2</sub> (1:1, 0.13 M) at -20°C was added I<sub>2</sub> (1.5 eq), KI (1.5 eq) then NaNO<sub>2</sub>(1.5 eq). The mixture was stirred at 20°C for 2 h then diluted with EtOAc and water. The organic extracts were washed with aqueous NaHCO<sub>3</sub>, brine, dried over MgSO<sub>4</sub>, filtered and concentrated. Flash chromatography (Hex:EtOAc ; 40:60) afforded the title compound as a white solid.

20


MC054PV

**Pyr\_5****3-bromo-5-phenylpyridine 1-oxide**

- 5 Prepared according to the procedure described for **Pyr\_2** using phenylboronic acid as starting material. Flash chromatography (Hex:EtOAc ; 30:70), stirring vigorously in EtOAc-Et<sub>2</sub>O-hexane then isolated by filtration.

**Pyr\_6****3-bromo-5-[3,5-(dichloro)phenyl]pyridine 1-oxide**

- 10  
15 Prepared according to the procedure described for **Pyr\_2** using 3,5-dichloro-phenylboronic acid as starting material. Flash chromatography (Hex:EtOAc ; 30:70), stirring vigorously in EtOAc-Et<sub>2</sub>O then isolated by filtration.


**Pyr\_7****3-bromo-5-[3,4-(methoxy)phenyl]pyridine 1-oxide**



MC054PV

Prepared according to the procedure described for Pyr\_2 using 3,4-dimethoxyphenylboronic acid as starting material. Flash chromatography (Hex:EtOAc ; 40:70 and then using toluene:acetone, 95:5) afforded the title compound.

5

**Pyr\_8****3-bromo-5-(5-methyl-1,2,4-oxadiazol-3-yl)pyridine 1-oxide**10 **Step 1 : 5-bromo-*N*'-hydroxypyridine-3-carboximidamide**

A mixture of 3-bromo-5-cyanopyridine (1.0 eq), hydroxylamine hydrochloride (1.5 eq.),  $K_2CO_3$  (2.1 eq) in ethanol (0.1 M) was stirred at 80°C for 3 days. The mixture was concentrated, diluted with EtOAc and aqueous  $NaHCO_3$ . The title compound was isolated by filtration as beige solid.

15

**Step 2 : 3-bromo-5-(5-methyl-1,2,4-oxadiazol-3-yl)pyridine**

To a solution of 5-bromo-*N*'-hydroxypyridine-3-carboximidamide (1.0 eq.) in pyridine was added acetyl chloride (2 eq). The reaction stirred at 100°C for 1 h and 2 eq of acetyl chloride was added. After 1 h, the mixture was cooled, diluted with EtOAc and aqueous  $NaHCO_3$ . The organic extracts were washed with aqueous  $NaHCO_3$ , brine, dried over  $Na_2SO_4$ , filtered and concentrated. Flash chromatography (Hex:EtOAc ; 70:30) afforded the title compound as a white solid.

20

25


**Step 3 : 3-bromo-5-(5-methyl-1,2,4-oxadiazol-3-yl)pyridine 1-oxide**

Prepared according to the procedure Oxid-2. Flash chromatography (EtOAc) afforded the title compound as a white solid.

MC054PV

## Pyr\_9

## 3-bromo-5-(5-methyl-1,3,4-oxadiazol-2-yl)pyridine 1-oxide



5

Step 1 : 3-bromo-5-(1*H*-tetraazol-5-yl)pyridine

To a mixture of 3-bromo-5-cyanopyridine (1.0 eq.) and triethylamine (1.3 eq in  
toluene (0.7 M) was added AcOH (1.3 eq). The reaction was stirred at 100°C for 12  
h, cooled, diluted with EtOAc and 1% AcOH solution. The organic extracts were  
washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated to afford the title  
compound as a brown solid.

15

## Step 2 : 3-bromo-5-(5-methyl-1,3,4-oxadiazol-2-yl)pyridine

A solution of 3-bromo-5-(5-methyl-1,3,4-oxadiazol-2-yl)pyridine in acetic anhydride  
(0.8M) was at 140°C for 2. The mixture was concentrated, diluted with EtOAc and  
aqueous NaHCO<sub>3</sub>. The organic extracts were washed with aqueous NaHCO<sub>3</sub>, brine,  
dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated. Flash chromatography (Hex:EtOAc ;  
70:30) afforded the title compound as a white solid.

20

## Step 3 : 3-bromo-5-(5-methyl-1,3,4-oxadiazol-2-yl)pyridine 1-oxide


Prepared according to the procedure Oxid-2. Flash chromatography (EtOAc:MeOH,  
100:0-95:5) afforded the title compound.

25

MC054PV

## Pyr\_10

## 2-(benzyloxy)-5-bromo-3-(3-methyl-1,2,4-oxadiazol-5-yl)pyridine



## 5 Step 1 : 2-chloro-3-(3-methyl-1,2,4-oxadiazol-5-yl)pyridine

A mixture of 2-chloro nicotinoyl chloride (1.0 eq.) and *N*<sup>2</sup>-hydroxy ethanimidamide (1.1 eq) in pyridine (0.6 M) was stirred at 100°C for 4 h and then concentrated. The residue was purified by flash chromatography (Hexane:EtOAc, 70:30) to afforded the title compound.

## Step 2 : 2-(benzyloxy)-3-(3-methyl-1,2,4-oxadiazol-5-yl)pyridine

To a solution of benzyl alcohol (1.5 eq) in DMF (0.4M) was added NaH (1.5 eq, 60%) followed by 2-chloro-3-(3-methyl-1,2,4-oxadiazol-5-yl)pyridine (1 eq) in DMF (0.5M) after 30 min. ). The reaction was stirred at 20°C for 1 h, diluted with Et<sub>2</sub>O and water. The organic extracts were washed with brine, dried over MgSO<sub>4</sub>, filtered and concentrated. Flash chromatography (Hex:EtOAc ; 80:20) afforded the title compound.

## 20 Step 3 : 3-(3-methyl-1,2,4-oxadiazol-5-yl)pyridin-2-ol

A solution of 2-(benzyloxy)-3-(3-methyl-1,2,4-oxadiazol-5-yl)pyridine in CH<sub>2</sub>Cl<sub>2</sub>-TFA (0.25M) was stirred 15 min at 20°C and then concentrated. The residu was diluted with EtOAc and aqueous NaHCO<sub>3</sub>. The organic extracts were washed with brine, dried over MgSO<sub>4</sub>, filtered and concentrated to afforded the title compound.

## Step 4 : 5-bromo-3-(3-methyl-1,2,4-oxadiazol-5-yl)pyridin-2-ol

MC054PV


A solution 3-(3-methyl-1,2,4-oxadiazol-5-yl)pyridin-2-ol in AcOH (0.2M) and bromine (1.2 eq) was stirred 18 h at 20°C and then concentrated. The residue was co-evaporated with benzene (2X) and used as such in the next step.

5 Step 5 : 2-(benzyloxy)-5-bromo-3-(3-methyl-1,2,4-oxadiazol-5-yl)pyridine

A solution 5-bromo-3-(3-methyl-1,2,4-oxadiazol-5-yl)pyridin-2-ol (1 eq) in benzene (0.05M) was added silver carbonate (1 eq) and benzyl bromide (1.2 eq). The reaction mixture was stirred 6 h at 20°C and then concentrated. The residue was purified by  
10 flash chromatography (Hexane:EtOAc, 90:10) to afforded the title compound.

Pyr\_11

3-bromo-5-(3-methyl-1,2,4-oxadiazol-5-yl)pyridine 1-oxide



15

Step 1 : 3-bromo-5-(3-methyl-1,2,4-oxadiazol-5-yl)pyridine


A mixture of 5-bromo nicotinic acid(1.0 eq.) and CDI (1.1 eq) in DMF (0.25M) was stirred at 20 °C for 5 min and then *N*<sup>2</sup>-hydroxy ethanimidamide (1.1 eq) in pyridine  
20 (0.6 M) was added. The reaction mixture was stirred at 110°C for 15 h, cooled to rt and poured into water. The title compound was isolated by filtration as a solid.

Step 2 : 3-bromo-5-(3-methyl-1,2,4-oxadiazol-5-yl)pyridine 1-oxide

25 Prepared according to the procedure Ox-2. The residu was stirred vigorously in EtOAc then isolated by filtration.

Pyr\_12

MC054PV

**5-bromo-2-[4-(methylsulfonyl)phenyl]pyridine 1-oxide****5** Step 1: 5-bromo-2-[4-(methylthio)phenyl]pyridine


Prepared according to the procedure **Coupling-2** using [4-(methylthio)phenyl]boronic acid and 2,5-dibromo pyridine afforded the title compound in a mixture of isomer containing 2-bromo-5-[4-(methylthio)phenyl]pyridine.

**10****Step 2: 5-bromo-2-[4-(methylsulfonyl)phenyl]pyridine**

Prepared by submitting the mixture of isomer to the procedure **Oxid-1**. Flash chromatography (DCM:EtOAc 95:5-50:50) afforded the title compound (less polar isomer).

**15****Step 3 : 5-bromo-2-[4-(methylsulfonyl)phenyl]pyridine 1-oxide**

Prepared according to the procedure **Oxid-2**. Flash chromatography (DCM:EtOAc 75:25-0:100) afforded the title compound.


**20****Pyr\_13****2-bromo-5-[4-(methylsulfonyl)phenyl]pyridine 1-oxide****25**

MC054PV

Prepared according to the procedure as described for **Pyr\_12** using the more polar isomer isolated from step 2.

**Pyr\_14**

5      **2-chloro-3-[4-(methylsulfonyl)phenyl]-5-(trifluoromethyl)pyridine**




Prepared according to the procedure as described in Bioorg. Med. Chem. Lett., 1999, 9, 1715, US Patent 5,861,419, 1999.

10

**Pyr\_15**

**3-bromo-1-(4-chlorophenyl)pyridin-2(1H)-one**



15      **Step 1 : 1-(4-chlorophenyl)pyridin-2(1H)-one**

Prepared according to the procedure as described in Chem.Pharm.Bull., 1997, 45, 719. Flash chromatography (Hex:EtOAc ; 70:30) afforded the title compound.

20      **Step 2 : 3-bromo-1-(4-chlorophenyl)pyridin-2(1H)-one**


A solution 1-(4-chlorophenyl)pyridin-2(1H)-one (1 eq) and tetrabutylammonium tribromide (1.2 eq) in chloroform (0.1M) was stirred 6 h at reflux. The reaction mixture was diluted with aqueous NaHCO<sub>3</sub> solution and aqueous Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> solution.

25      The organic extracts were washed with brine, dried over MgSO<sub>4</sub>, filtered and

concentrated. The residue was purified by flash chromatography (Hexane:EtOAc, 80:20) to afford the title compound.

**Pyr\_16**

5      3-bromo-5-(2-methyl-2H-tetraazol-5-yl)pyridine 1-oxide



**Step 1: 3-bromo-5-(2H-tetraazol-5-yl)pyridine**

10      A solution of 3-bromo-5-cyanopyridine (1 eq), sodium azide (1.2 eq), triethylamine (1.2 eq) and AcOH (1.2 eq) was stirred at 100C for 14h. The reaction mixture was cooled to rt, diluted with aqueous acetic acid solution and extracted with EtOAc. The organic extracts were washed with brine, dried over MgSO<sub>4</sub>, filtered and concentrated to afford the title compound.

15

**Step 2: 3-bromo-5-(2-methyl-2H-tetraazol-5-yl)pyridine**


To a solution of tetrazole from step 1 in CH<sub>2</sub>Cl<sub>2</sub>-MeOH (5:1, 0.1M) was added CH<sub>2</sub>N<sub>2</sub> (in ether) until reaction completed by TLC. The reaction mixture was concentrated to  
20      afford the title compound.

**Step 3: 3-bromo-5-(2-methyl-2H-tetraazol-5-yl)pyridine 1-oxide**

Prepared according to the procedure Oxid-2. Flash chromatography (CH<sub>2</sub>Cl<sub>2</sub>-MeOH, 98:2) afforded the title compound.  
25

MC054PV

## Het\_1

5-bromo-*N*-isopropyl-4-[4-(methylsulfonyl)phenyl]thiophene-2-carboxamide

## 5 Step 1 : 3-[4-(methylthio)phenyl]thiophene

Prepared according to the procedure **Coupling-1** using [4-(methylthio)phenyl]boronic acid and 3-bromothiophene as starting material. Flash chromatography ( $\text{CH}_2\text{Cl}_2$ ) afforded the title compound.

10

## Step 2 : 2-bromo-3-[4-(methylthio)phenyl]thiophene

To a solution of thiophene from step 1 (1.0 eq) in  $\text{CH}_2\text{Cl}_2/\text{AcOH}$  (1:1, 0.1M) was added NBS (1.05 eq). The mixture was stirred 1.5 h at r.t., diluted with EtOAc and aqueous  $\text{NaHCO}_3$ . The organic extracts were washed with water, brine, dried over  $\text{MgSO}_4$ , filtered and concentrated. Flash chromatography (Hexane:EtOAc, 80:20) afforded the title compound.

15

Step 3 : 5-bromo-*N*-isopropyl-4-[4-(methylthio)phenyl]thiophene-2-carboxamide

20

To a solution of diisopropylamine (1.05 eq) in THF at  $-78^\circ\text{C}$  was added BuLi (1.6M, Hexane, 1.05 eq) dropwise. The mixture was stirred 10 min at  $0^\circ\text{C}$ , cooled to  $-78^\circ\text{C}$  and thiophene from step 2 added. After 30 min at  $-50^\circ\text{C}$ , isopropyl isocyanate was added. The mixture was stirred 1 h at rt, diluted with EtOAc and aqueous  $\text{NaHCO}_3$ . The organic extracts were washed with water, brine, dried over  $\text{MgSO}_4$ , filtered and concentrated. Flash chromatography ( $\text{CH}_2\text{Cl}_2$ ) afforded the title compound.

25




MC054PV

Step 4 : 5-bromo-*N*-isopropyl-4-[4-(methylsulfonyl)phenyl]thiophene-2-carboxamide

Prepared according to the procedure **Oxid-1**. Flash chromatography (CH<sub>2</sub>Cl<sub>2</sub>: EtOAc, 90:10) afforded the title compound.


5

**Phe\_1****2-bromo-4'-(methylthio)-1,1'-biphenyl**

- 10 Prepared according to the general procedure **Coupling-1** using 4-methylthiobenzene boronic acid and 1-bromo-2-iodobenzene. Flash chromatography (Hexane:EtOAc, 100:0-95:5) afforded the title compound as an oil.


**Phe\_2**

15

**2-(2-bromophenyl)-5-methyl-1,3,4-oxadiazole**

- 20 A mixture of 5-(2-bromophenyl)-2*H*-tetraazole (1.0 eq) in acetic anhydride (1M) was heated at 120°C for 2 h and then concentrated. Flash chromatography (Hexane:EtOAc, 70:30) afforded the title compound as a white solid.

MC054PV

**Phe\_3****methyl 6-iodo-4'-(methylsulfonyl)-1,1'-biphenyl-3-carboxylate****5 Step 1 : methyl 6-amino-4'-(methylthio)-1,1'-biphenyl-3-carboxylate**

Prepared according to the general procedure **Coupling-1** (CsF, DME, 20°C) using [4-(methylthio)phenyl]boronic acid and methyl 4-amino-3-iodobenzoate as starting material. Flash chromatography (Hexane:EtOAc, 70:30) afforded the title compound  
**10** as a white solid.

**Step 2 : methyl 6-amino-4'-(methylsulfonyl)-1,1'-biphenyl-3-carboxylate**

Prepared according to the general procedure **Oxid-1**. Flash chromatography  
**15** (Hexane:EtOAc, 50:50) afforded the title compound.


**Step 3 : methyl 6-iodo-4'-(methylsulfonyl)-1,1'-biphenyl-3-carboxylate**

To the amino from step 2 (1.0 eq) in CH<sub>2</sub>Cl<sub>2</sub>-AcOH (1:1) at -20°C was added iodine  
**20** (1.5 eq), KI (1.5 eq) followed by NaNO<sub>2</sub> (1.2 eq). The reaction mixture was stirred at r.t. for 1 h, diluted with EtOAc, a solution of Na<sub>2</sub>CO<sub>3</sub> and a solution Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub>. The organic extracts were washed with water, brine, dried over MgSO<sub>4</sub>, filtered and concentrated. Flash chromatography (Hexane:EtOAc, 30:20-20:30) afforded the title  
**25** compound as a pale yellow solid.

MC054PV

## Phe\_4

2-[5-(2-bromophenyl)-1,3-thiazol-2-yl]propan-2-ol



## 5 Step 1 : 2-[5-(tributylstannyl)-1,3-thiazol-2-yl]propan-2-ol

To a solution of thiazole (1.0 eq) in THF (0.2M) at  $-78^{\circ}\text{C}$  was added nBuLi (1.6M, 1.1 eq). After 1 h at  $-78^{\circ}\text{C}$ , acetone (1.0 eq) was added and the mixture stirred 30 min at  $-78^{\circ}\text{C}$ , 5 min  $-50^{\circ}\text{C}$  then cooled back to  $-78^{\circ}\text{C}$  to add a second equivalent of nBuLi (1.6M, 1.1 eq). After 1.5 h at  $-78^{\circ}\text{C}$ , tri-n-butyltin chloride (1.2 eq) was added and the mixture stirred 12 h at  $-78^{\circ}\text{C}$ . The reaction mixture was diluted with  $\text{Et}_2\text{O}$  and a solution of  $\text{NH}_4\text{Cl}$ . The organic extracts were washed with water, brine, dried over  $\text{MgSO}_4$ , filtered and concentrated. Flash chromatography (Hexane:EtOAc, 90:10) afforded the title compound.


15

## Step 2: 2-[5-(2-bromophenyl)-1,3-thiazol-2-yl]propan-2-ol

A solution of stannane from step 1 (1.0 eq), 1-bromo-2-iodo-benzene (2.1 eq), CuI (0.2 eq),  $\text{PdCl}_2(\text{dppf})$  (0.1 eq) in DMF was heated 1 h at  $80^{\circ}\text{C}$ . The reaction mixture was diluted with EtOAc and a solution of  $\text{NH}_4\text{Cl}$ . The organic extracts were washed with water, brine, dried over  $\text{MgSO}_4$ , filtered and concentrated. Flash chromatography (Hexane:EtOAc, 80:20) afforded the title compound.

20

MC054PV

**Phe\_5****4-(3-bromophenyl)pyridine 1-oxide****5 Step 1 : 4-(3-bromophenyl)pyridine**

Prepared according to the general procedure **Coupling-1** using 4-pyridinylboronic acid and 1,3-dibromobenzene. Flash chromatography (Hexane:EtOAc, 70:30) afforded the title compound.

10


**Step 2 : 4-(3-bromophenyl)pyridine 1-oxide**

Prepared according to the general procedure **Oxid-2**. Flash chromatography (EtOAc:MeOH, 100:0 to 95:5) afforded the title compound.

15

**EXAMPLE 1**

**8-(3-{2-(3-chlorophenyl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-5-yl}phenyl)-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline**



20

MC054PV

Step 1 : diphenyl anilino[4-(methylthio)phenyl]methylphosphonate

To a solution of 4-methylthiobenzaldehyde (1.0 eq) dissolved in hot acetonitrile was added aniline (1.2 eq) and diphenyl phosphite (1.6 eq). The reaction mixture was stirred at 20°C for 18 h and then concentrated. The residue can be purified by flash chromatography if necessary.

Step 2 : 2-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-1-[4-(methylthio)phenyl]ethanone

10

To a solution of phosphonate from step 1 (1.2 eq) dissolved in DMF (0.2M) was added cesium carbonate (1.5 eq), **Quinoline\_2** (1.0 eq) and isopropanol (20%, v/v).

The reaction mixture was stirred at 70°C for 30 min (or 20°C for 18 h), cooled to rt diluted with HCl (6N), stirred for 2 h and extracted with EtOAc. The organic extracts were washed with brine, dried over MgSO<sub>4</sub>, filtered and concentrated. The residue was purified by flash chromatography (Hexane:EtOAc, 50:50) to afforded the title compound.

Step 3 : 2-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-1-[4-(methylsulfonyl)phenyl]ethanone

20

Prepared according to the procedure **Oxid-1**. Flash chromatography (Hexane:EtOAc) afforded the title compound.

Step 4 : 2-bromo-2-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-1-[4-(methylsulfonyl)phenyl]ethanone.

25

A solution of ketone from step 3 (1.0 eq) in chloroform (0.2M) with polymer-supported tribromide (1 mmol/g, 1.2 eq) was heated at reflux for 1-2 h, filtered and concentrated to afforded the title compound.

30

Step 5 : 8-(3-{2-(3-chlorophenyl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-5-yl}phenyl)-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline

MC054PV

To a solution of the alpha-bromoketone from step 4 (1.0 eq) in DMF (0.1M) was added 3-chlorobenzenecarbothioamide (1.0 eq). The mixture was heated at 40-120°C for 1-3 h, cooled, diluted with water, saturated NaHCO<sub>3</sub> solution and ether. The organic extracts were washed with brine, dried over MgSO<sub>4</sub>, filtered and concentrated. The residue was purified by flash chromatography (Hexane:EtOAc) to afford the title compound.

<sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>): δ 8.85 (dd, 1H), 8.46 (dd, 1H), 8.22 (d, 1H), 8.04 (s, 1H), 7.95-7.90 (m, 6H), 7.73 (m, 2H), 7.55 (m, 4H), 7.48 (d, 1H), 3.28 (s, 3H), 2.74 (s, 3H), 1.85 (s, 6H).

Following examples (Table 1) were prepared according to the procedure described for EXAMPLE 1. Non-commercial thioamide used in step 5 of example 1, were prepared according to general procedure Thio-1.




Table 1:

Ex #:	R group name	ESI-LRMS (M + 1) <sup>+</sup>
2	Amino	709.8
3	2-biphenyl	636.5
4	3-biphenyl	636.6
5	N-(tert-butoxycarbonyl)-4-phenylpyrrolidin-3-yl	729.7
6	N-(tert-butoxycarbonyl)azetidin-3-yl	639.6
7	N-(tert-butoxycarbonyl)pyrrolidin-3-yl	653.9
8	3-chloro-4-fluorophenyl	611.3
9	4-chlorophenoxymethyl	624.4
10	2-chlorophenyl	594.5

MC054PV


11	4-chlorophenyl	593.3
12	Ethoxycarbonyl	635.3
13	furan-2-yl	550.5
14	furan-3-yl	550.4
15	imidazol-2-yl	550.4
16	indan-1-yl	600.6
17	indan-2-yl	600.6
18	1H-indol-2-yl	599.5
19	1H-indol-3-yl	599.6
20	1H-indol-4-yl	599.4
21	1H-indol-5-yl	599.5
22	1H-indol-6-yl	599.6
23	1H-indol-7-yl	599.5
24	isoquinolin-1-yl	611.5
25	isoquinolin-4-yl	611.5
26	isoquinolin-5-yl	611.5
27	isoquinolin-8-yl	690.5
28	isoxazol-3-yl	551.5
29	3-methoxycarbonylphenyl	697.2
30	4-methoxycarbonylphenyl	697.2
31	Methyl	577.3
32	1-methyl-1H-pyrazol-3-yl	564.5
33	1-methyl-1H-pyrazol-4-yl	564.6
34	1-methyl-1H-pyrazol-5-yl	564.5
35	2-methylphenyl	653.3
36	3-methylphenyl	653.3
37	4-methylphenyl	574.6
38	2-methylpyridin-5-yl	575.5
39	Methylsulfonylmethyl	576.5
40	2-methylsulfonylphenyl	717.3
41	3-methylsulfonylphenyl	717.3
42	4-methylsulfonylphenyl	717.3
43	morpholin-4-ylmethyl	575.5

44	Phenyl	639.3
45	Pyrazinyl	562.4
46	1H-pyrazol-3-yl	550.4
47	pyridin-2-yl	561.5
48	pyridin-3-yl	561.5
49	pyridin-4-yl	561.5
50	3-pyridinylmethyl	575.6
51	pyrimidin-2-yl	562.4
52	pyrimidin-4-yl	562.4
53	pyrimidin-5-yl	562.4
54	quinolin-4-yl	611.5
55	quinolin-5-yl	611.5
56	quinolin-8-yl	611.5
57	5,6,7,8-tetrahydro-1,8-naphthyridin-2-yl	695.6
58	6,7,8,9-tetrahydro-5H-benzo[a][7]annulen-5-yl	628.6
59	6,7,8,9-tetrahydro-5H-benzo[a][7]annulen-6-yl	628.6
60	tetrahydrofuran-2-yl	554.5
61	1,2,3,4-tetrahydronaphthalen-1-yl	615.7
62	1,2,3,4-tetrahydronaphthalen-2-yl	615.6
63	1,3-thiazol-2-yl	567.3
64	1,3-thiazol-5-yl	567.4
65	thiophen-2-yl	566.2
66	thiophen-3-yl	566.3



## EXAMPLE 67

6-[1-methyl-1-(methanesulfonyl)ethyl]-8-{3-[4-[4-(methanesulfonyl)phenyl]-2-(1-oxido-2,3-dihydropyridin-4-yl)-1,3-thiazol-5-yl]phenyl}quinoline




5

Prepared according to the general procedure Oxid-2 using EXAMPLE 49 as starting material.

<sup>1</sup>H NMR (500 MHz, acetone-*d*<sub>6</sub>): δ 8.88 (dd, 1H), 8.45 (dd, 1H), 8.28-8.26 (m, 3H), 8.10 (d, 1H), 8.04 (d, 2H), 8.01 (d, 2H), 7.96 (d, 2H), 7.86 (s, 1H), 7.83 (d, 1H), 7.61 (t, 1H), 7.58-7.55 (m, 2H), 3.14 (s, 3H), 2.69 (s, 3H), 1.96 (s, 6H).  
ESI-LRMS (M + 1) 656.4

## EXAMPLE 68

6-[1-methyl-1-(methanesulfonyl)ethyl]-8-{3-[4-[4-(methanesulfonyl)phenyl]-2-(1-oxido-2,3-dihydropyridin-3-yl)-1,3-thiazol-5-yl]phenyl}quinoline



20 Prepared according to the general procedure Oxid-2 using EXAMPLE 48 as starting material.

MC054PV


<sup>1</sup>H NMR (500 MHz, acetone-*d*<sub>6</sub>): δ 8.88 (dd, 1H), 8.79 (s, 1H), 8.44 (dd, 1H), 8.28 (d, 1H), 8.25 (d, 1H), 8.10 (d, 1H), 8.03 (d, 2H), 7.97 (d, 2H), 7.89-7.84 (m, 3H), 7.63 (t, 1H), 7.58-7.47 (m, 3H), 3.15 (s, 3H), 2.70 (s, 3H), 1.96 (s, 6H).

ESI-LRMS (M + 1) 656.3

5

## EXAMPLE 69

2-(3-{5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-2-yl}phenyl)propan-2-ol



- 10 Prepared according to the general procedure Cer-1 using methyl magnesium bromide and EXAMPLE 29 as starting material.

<sup>1</sup>H NMR (500 MHz, acetone-*d*<sub>6</sub>): δ 8.89 (dd, 1H), 8.44 (dd, 1H), 8.28 (dd, 2H), 8.09 (d, 1H), 8.01 (d, 2H), 7.95 (d, 2H), 7.92 (d, 1H), 7.86 (s, 1H), 7.82 (d, 1H), 7.66 (d, 1H), 7.60 (t, 1H), 7.57-7.54 (m, 2H), 7.48 (t, 1H), 4.28 (s, 0.6H), 3.13 (s, 3H), 2.69 (s, 3H), 1.96 (s, 6H), 1.58 (s, 6H).


15

ESI-LRMS (M + 1) 617.4

MC054PV

## EXAMPLE 70

3-{5-[3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-2-yl}benzoic acid



5

10

15

To a solution of **EXAMPLE 29** in THF/MeOH/H<sub>2</sub>O (0.05M) was added LiOH (2N, 5 eq). The reaction mixture was stirred at rt for 2h, acidified with AcOH, diluted with water and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic extracts were washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated. The title compound was isolated by filtration from acetone/ether as a solid.


<sup>1</sup>H NMR (500 MHz, acetone-*d*<sub>6</sub>): δ 8.89 (dd, 1H), 8.72 (s, 1H), 8.44 (dd, 1H), 8.31 (d, 1H), 8.27 (d, 1H), 8.17 (d, 1H), 8.10 (d, 1H), 8.03 (d, 2H), 7.96 (d, 2H), 7.88 (s, 1H), 7.83 (d, 1H), 7.69 (t, 1H), 7.61 (t, 1H), 7.58-7.54 (m, 2H), 3.14 (s, 3H), 2.69 (s, 3H), 1.96 (s, 6H).

ESI-LRMS (M + 1) 683.3

MC054PV

## EXAMPLE 71

2-{5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-2-yl}propan-2-ol




Prepared according to the general procedure Cer-1 using methyl magnesium bromide and EXAMPLE 12 as starting material.

<sup>1</sup>H NMR (500 MHz, acetone-*d*<sub>6</sub>): δ 8.88 (dd, 1H), 8.45 (dd, 1H), 8.27 (d, 1H), 8.09 (d, 1H), 7.90 (m, 4H), 7.77 (m, 2H), 7.56 (m, 2H), 7.47 (m, 1H), 5.26 (s, OH), 3.10 (s, 3H), 2.70 (s, 3H), 1.95 (s, 6H), 1.70 (s, 6H).

## EXAMPLE 72

6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{2-(3-methyl-1,2,4-oxadiazol-5-yl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-5-yl}phenyl)quinoline



MC054PV

Step 1: 5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazole-2-carboxylic acid

To a solution of **EXAMPLE 12** in THF/MeOH/H<sub>2</sub>O (0.05M) was added LiOH (2N, 2 eq). The reaction mixture was stirred at rt for 2h, acidified with HCO<sub>2</sub>H, diluted with water and extracted with EtOAc. The organic extracts were washed with brine and concentrated to afford the title compound as a solid.


Step 2: 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{2-(3-methyl-1,2,4-oxadiazol-5-yl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-5-yl}phenyl)quinoline

To a solution of acid from step 1 in DMF (0.05M) was added CDI (1.1 eq) and, after 5 min, acetamidoxime (1.1 eq). The reaction mixture was stirred at rt for 1h and 18h at 110°C. The reaction mixture was cooled to rt and poured into water. The residue was purified by flash chromatography (toluene:EtOAc, 45:55) to afford the title compound and decarboxylated analog **EXAMPLE 73** (more polar product).

<sup>1</sup>H NMR (500 MHz, acetone-*d*<sub>6</sub>): δ 8.88 (dd, 1H), 8.47 (dd, 1H), 8.30 (d, 1H), 8.12 (d, 1H), 8.01 (s, 4H), 7.90 (m, 2H), 7.60 (m, 3H), 3.18 (s, 3H), 2.70 (s, 3H), 2.52 (s, 3H), 1.95 (s, 6H).

#### EXAMPLE 73

6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-5-yl}phenyl)quinoline



Prepared according to the procedure described for **EXAMPLE 72**.


MC054PV

<sup>1</sup>H NMR (500 MHz, acetone-*d*<sub>6</sub>): δ 9.15 (s, 1H), 8.88 (dd, 1H), 8.43 (dd, 1H), 8.25 (d, 1H), 8.08 (d, 1H), 7.92 (m, 4H), 7.80 (m, 2H), 7.59 (t, 1H), 7.55 (dd, 1H), 7.50 (m, 1H), 3.12 (s, 3H), 2.70 (s, 3H), 1.95 (s, 6H).

5

## EXAMPLE 74

*N*-cyclopropyl-5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazole-2-carboxamide



10


To a solution of **EXAMPLE 12** in CH<sub>2</sub>Cl<sub>2</sub> (0.05M) was added cyclopropylamine (4 eq) and trimethylaluminum (2M, Hexanes, 4 eq). The reaction mixture was stirred at rt for 18h. The reaction mixture was diluted slowly with a saturated NH<sub>4</sub>Cl solution, a  
 15 1M sodium potassium tartrate solution and EtOAc. The mixture was extracted with EtOAc, the organic extracts were washed with brine, dried over MgSO<sub>4</sub>, filtered and concentrated. The residue was purified by flash chromatography (EtOAc) to afford the title compound.

<sup>1</sup>H NMR (500 MHz, acetone-*d*<sub>6</sub>): δ 8.86 (dd, 1H), 8.43 (dd, 1H), 8.31 (d, 1H), 8.26 (d, 1H), 8.07 (d, 1H), 7.91 (s, 4H), 7.83 (m, 2H), 7.55 (m, 3H), 3.12 (s, 3H), 2.95 (m, 20 1H), 2.69 (s, 3H), 1.95 (s, 6H), 0.78 (m, 2H), 0.73 (m, 2H).

MC054PV

## EXAMPLE 75

6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{2-(6-methyl-1-oxidopyridin-3-yl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-5-yl}phenyl)quinoline



5

Prepared according to the general procedure Oxid-2 using EXAMPLE 38 as starting material.


<sup>1</sup>H NMR (500 MHz, acetone-*d*<sub>6</sub>): δ 8.87 (m, 2H), 8.44 (dd, 1H), 8.26 (d, 1H), 8.09 (d, 1H), 8.0-7.9 (m, 2H), 8.0-7.9 (m, 5H), 7.67-7.5 (m, 4H), 3.13 (s, 3H), 2.7 (s, 3H), 2.45 (s, 3H), 1.95 (s, 6H).

10

## EXAMPLE 76

2-[4-(4-chlorophenyl)-5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-1,3-thiazol-2-yl]propan-2-ol

15



MC054PV

Step 1: ethyl 4-(4-chlorophenyl)-5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl] quinolin-8-yl}phenyl)-1,3-thiazole-2-carboxylate

Prepared according to the procedure described for **EXAMPLE 1** using 4-chloro benzaldehyde in step 1 and ethyl thiooxamate in step 5. Flash chromatography (Hexane:EtOAc, 50:50-20:80) afforded the title compound.


Step 2 : 2-[4-(4-chlorophenyl)-5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-1,3-thiazol-2-yl]propan-2-ol

Prepared according to the general procedure Cer-1 using methyl magnesium bromide. Flash chromatography (Hexane:EtOAc, 35:65) afforded the title compound as a white foam.

<sup>1</sup>H NMR (500 MHz, acetone-*d*<sub>6</sub>): δ 8.85 (dd, 1H), 8.40 (dd, 1H), 8.25 (d, 1H), 8.05 (s, 1H), 7.75 (m, 2H), 7.65 (d, 3H), 7.35 (d, 2H), 7.0 (m, 2H), 5.10 (s, OH), 2.66 (s, 3H), 1.95 (s, 6H), 1.7 (s, 6H).

#### EXAMPLE 77

6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{2-methyl-4-[4-(methylsulfonyl)phenyl]-1,3-oxazol-5-yl}phenyl)quinoline



Prepared according to the procedure described for **EXAMPLE 1** using acetamide at 120°C in step 5. Flash chromatography (CH<sub>2</sub>Cl<sub>2</sub>: EtOAc, 80:20) afforded the title compound.




MC054PV

<sup>1</sup>H NMR (500 MHz, acetone-*d*<sub>6</sub>): δ 8.94 (dd, 1H), 8.45 (dd, 1H), 8.27 (d, 1H), 8.12 (d, 1H), 8.03 (d, 2H), 7.98 (m, 3H), 7.80 (d, 1H), 7.70 (m, 1H), 7.62 (m, 1H), 7.57 (m, 1H), 3.12 (s, 3H), 2.7 (s, 3H), 2.52 (s, 3H), 1.95 (s, 6H).

EXAMPLE 78

2-{4-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-5-[4-(methylsulfonyl)phenyl]-1,3-thiazol-2-yl}propan-2-ol



Step 1: ethyl 4-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-5-[4-(methylthio)phenyl]-1,3-thiazole-2-carboxylate

Prepared according to the procedure described for **EXAMPLE 1** using **Quinoline\_2** in step 1, 4-methylthiobenzaldehyde in step 2 and ethyl thiooxamate in step 5.

Step 2: 2-{4-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-5-[4-(methylthio)phenyl]-1,3-thiazol-2-yl}propan-2-ol

Prepared according to the general procedure **Cer-1** using methyl magnesium bromide. Flash chromatography (CH<sub>2</sub>Cl<sub>2</sub>:MeOH, 98:2) afforded the title compound.

Step 3: 2-{4-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-5-[4-(methylsulfonyl)phenyl]-1,3-thiazol-2-yl}propan-2-ol


MC054PV

Prepared according to the general procedure **Oxid-1**. Flash chromatography (toluene:acetone, 70:30) afforded the title compound.

<sup>1</sup>H NMR (500 MHz, acetone-*d*<sub>6</sub>): δ 8.89 (dd, 1H), 8.43 (dd, 1H), 8.26 (d, 1H), 8.05 (d, 1H), 7.96 (d, 2H), 7.87 (t, 1H), 7.73 (d, 2H), 7.72 (d, 1H), 7.63 (d, 1H), 7.55 (dd, 1H), 7.50 (t, 1H), 5.20 (s, OH), 3.13 (s, 3H), 2.71 (s, 3H), 1.97 (s, 6H), 1.71 (s, 6H).

#### EXAMPLE 79

1,1,1-trifluoro-*N*-{5-(3-(6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl)phenyl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-2-yl}methanesulfonamide




To a suspension of **EXAMPLE 2** in CH<sub>2</sub>Cl<sub>2</sub> at -78°C was added triethylamine (4 eq) and Tf<sub>2</sub>O (2.2 eq). The reaction was stirred at -78°C 15 min then warmed to 20°C for 1 h, diluted with NaHCO<sub>3</sub> solution and CH<sub>2</sub>Cl<sub>2</sub> and stirred 15 min. The organic extracts were washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated. Flash chromatography (EtOAc) afforded the title compound.

<sup>1</sup>H NMR (500 MHz, acetone-*d*<sub>6</sub>): δ 8.89 (dd, 1H), 8.43 (dd, 1H), 8.27 (d, 1H), 8.06 (d, 1H), 7.89 (d, 2H), 7.87 (d, 2H), 7.73-7.70 (m, 2H), 7.56 (dd, 1H), 7.52 (t, 1H), 7.43 (dd, 1H), 3.12 (s, 3H), 2.70 (s, 3H), 1.98 (s, 6H).

MC054PV

## EXAMPLE 80

2-[5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-4-pyridin-3-yl-1,3-thiazol-2-yl]propan-2-ol



5

Step 1: ethyl 4-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-5-pyridin-3-yl-1,3-thiazole-2-carboxylate

10

Prepared according to the procedure described for **EXAMPLE 1** using 3-pyridinecarboxaldehyde in step 1 and ethyl thiooxamate in step 5. Flash chromatography (Hexane:EtOAc, 50:50-0:100) afforded the title compound.

15

Step 2: 2-[4-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-5-pyridin-3-yl-1,3-thiazol-2-yl]propan-2-ol

Prepared according to the general procedure **Cer-1** using methyl magnesium bromide. Flash chromatography (CH<sub>2</sub>Cl<sub>2</sub>:MeOH, 96:4) afforded the title compound.


20

<sup>1</sup>H NMR (500 MHz, acetone-*d*<sub>6</sub>): δ 8.86-8.85 (m, 2H), 8.49 (dd, 1H), 8.39 (dd, 1H), 8.24 (d, 1H), 8.05 (d, 1H), 7.97 (dt, 1H), 7.78-7.75 (m, 2H), 7.54-7.50 (m, 2H), 7.45 (dt, 1H), 7.33 (dd, 1H), 5.26 (br s, OH), 2.67 (s, 3H), 1.94 (s, 6H), 1.69 (s, 6H).

MC054PV

## EXAMPLE 81

2-[5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-4-(1-  
 oxidopyridin-3-yl)-1,3-thiazol-2-yl]propan-2-ol



5

Prepared according to the general procedure Oxid-2 using EXAMPLE 80 as starting material. Flash chromatography ( $\text{CH}_2\text{Cl}_2$ :MeOH, 95:5) afforded the title compound.


$^1\text{H}$  NMR (500 MHz, acetone- $d_6$ ):  $\delta$  8.89 (dd, 1H), 8.43 (d, 1H), 8.26 (s, 1H), 8.00 (s, 1H), 7.82 (s, 1H), 7.72 (d, 1H), 7.68 (d, 1H), 7.58 (dd, 1H), 7.50-7.43 (m, 5H), 5.12

10 (s, OH), 2.68 (s, 3H), 1.98 (s, 6H), 1.70 (s, 6H).

## EXAMPLE 82

1-(4-chlorophenyl)-1-{4-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-  
 yl}phenyl)-5-[4-(methylsulfonyl)phenyl]-1,3-thiazol-2-yl}ethanol

15



MC054PV

Step 1: ethyl 4-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-5-[4-(methylthio)phenyl]-1,3-thiazole-2-carboxylate

Prepared according to the procedure described for **EXAMPLE 1** using **Quinoline\_2** in step 1, 4-methylthiobenzaldehyde in step 2 and ethyl thiooxamate in step 5. (step 3 is omitted).

Step 2: *N*-methoxy-*N*-methyl-4-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-5-[4-(methylthio)phenyl]-1,3-thiazole-2-carboxamide

To a solution of methoxymethylamine in THF (0.08M) at  $-78^{\circ}\text{C}$  was added methylmagnesium iodide (3M, ether, 1 eq) dropwise over 1h. The reaction mixture was stirred at  $-78^{\circ}\text{C}$  for 10 min, a solution of ester from step 1 was added and the reaction mixture was stirred at  $-78^{\circ}\text{C}$  for 2h. The mixture was quenched with AcOH, diluted with  $\text{NH}_4\text{Cl}$  solution and EtOAc. The organic extracts were washed with brine, dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated. The residue was purified by flash chromatography (toluene:acetone, 80:20) to afford the title compound.

Step 3: 1-{4-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-5-[4-(methylthio)phenyl]-1,3-thiazol-2-yl}ethanone

Prepared according to the general procedure **Cer-1** using methylmagnesium iodide and amide from step 2.

Step 4: 1-(4-chlorophenyl)-1-{4-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-5-[4-(methylthio)phenyl]-1,3-thiazol-2-yl}ethanol

Prepared according to the general procedure **Cer-1** using 4-chlorophenylmagnesium bromide and methylketone from step 3. Flash chromatography ( $\text{CH}_2\text{Cl}_2$ :MeOH, 96:4) afforded the title compound.

MC054PV

Step 5: 1-(4-chlorophenyl)-1-{4-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-5-[4-(methylsulfonyl)phenyl]-1,3-thiazol-2-yl}ethanol

Prepared according to the general procedure **Oxid-1** using sulfide from step 4 as starting material. Flash chromatography ( $\text{CH}_2\text{Cl}_2$ :MeOH, 99:1) afforded the title compound.

$^1\text{H}$  NMR (500 MHz, acetone- $d_6$ ):  $\delta$  8.90 (dd, 1H), 8.44 (dd, 1H), 8.26 (d, 1H), 8.07 (d, 1H), 7.95 (dd, 2H), 7.90 (t, 1H), 7.83 (dd, 2H), 7.76 (dd, 1H), 7.71 (dd, 2H), 7.62 (dt, 1H), 7.56 (dd, 1H), 7.50 (t, 1H), 7.40 (dd, 2H), 6.05 (s, OH), 3.12 (s, 3H), 2.72 (s, 3H), 2.11 (s, 3H), 1.98 (s, 6H).

The following compounds (**Table 2**) were prepared according to the procedure described previously (Ex. 1-80). Indicated is their respective  $(M + 1)^+$  value obtained from a low resolution mass spectrometer under electron-spray ionization conditions.

**Table 2:**


EX.	Chemical name	ESI-LRMS (M + 1) <sup>+</sup>
83	6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{2-[3-(methylsulfonyl)phenyl]-4-phenyl-1,3-thiazol-5-yl}phenyl)quinoline	639.3
84	2-[4-(3-chlorophenyl)-5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-1,3-thiazol-2-yl]propan-2-ol	577.2
85	2-[4-(4-fluorophenyl)-5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-1,3-thiazol-2-yl]propan-2-ol	561.2
86	8-{3-[4-(4-chlorophenyl)-2-quinolin-5-yl]-1,3-thiazol-5-yl}phenyl-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline	646.2
87	2-{3-[4-(3-chlorophenyl)-5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-1,3-thiazol-2-yl]phenyl}propan-2-ol	653.5
88	2-{3-[4-(3-chloro-4-fluorophenyl)-5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-1,3-thiazol-2-yl]phenyl}propan-2-ol	670.9

MC054PV

89	2-{3-[4-[3,4-bis(difluoromethoxy)phenyl]-5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-1,3-thiazol-2-yl]phenyl}propan-2-ol	751.3
90	<i>N</i> -(5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-2-yl)acetamide	619.7
91	<i>N</i> -(5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-2-yl)pyridin-4-amine	655.4
92	2-[5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-4-pyridin-4-yl-1,3-thiazol-2-yl]propan-2-ol	543.9
93	2-[5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-4-(1-oxidopyridin-4-yl)-1,3-thiazol-2-yl]propan-2-ol	560.7
94	2-[5-(4-chlorophenyl)-4-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-1,3-thiazol-2-yl]propan-2-ol	577.2
95	2-{3-[4-(4-chlorophenyl)-5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-1,3-thiazol-2-yl]phenyl}propan-2-ol	653.6
96	6-[1-methyl-1-(methylsulfonyl)ethyl]-8-{3-[5-(1 <i>H</i> -tetrazol-5-yl)pyridin-3-yl]phenyl}quinoline	471.1

#### EXAMPLE 97

6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{3-[4-(methylsulfonyl)phenyl]thien-2-yl}phenyl)quinoline



5

Step 1: 2-bromo-3-[4-(methylthio)phenyl]thiophene

MC054PV

Prepared according to the general procedure **Coupling-1** using 4-methylthiobenzeneboronic acid and 3-bromothiophene. Flash chromatography ( $\text{CH}_2\text{Cl}_2$ ) afforded the title compound.

5 Step 2: 4-(2-bromothiophen-3-yl)phenyl methyl sulfone

Prepared according to the general procedure **Oxid-1** using sulfide from step 1 as starting material.

10 Step 3: 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{3-[4-(methylsulfonyl)phenyl]thien-2-yl}phenyl)quinoline.

Prepared according to the general procedure **Coupling-1** using **Quinoline\_4** and 3-bromothiophene from step 2 as starting material. Flash chromatography


15 (toluene:EtOAc ; 70:30) afforded the title compound.

$^1\text{H}$  NMR (500 MHz, acetone- $d_6$ ):  $\delta$  8.88 (dd, 1H), 8.45 (dd, 1H), 8.25 (d, 1H), 8.03 (d, 1H), 7.89 (d, 2H), 7.70 (m, 2H), 7.62 (m, 3H), 7.55 (dd, 1H), 7.47 (t, 1H), 7.37 (m, 1H), 7.32 (d, 1H), 3.08 (s, 3H), 2.70 (s, 3H), 1.95 (s, 6H).

20

EXAMPLE 98

1-(3'-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}-1,1':2',1''-terphenyl-4-yl)ethanone



25



MC054PV

Step 1 : 8-(2'-bromo-1,1'-biphenyl-3-yl)-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline

Prepared according to the general procedure **Coupling-1** using **Quinoline\_4** and 1,2-dibromobenzene as starting material. Flash chromatography (Hex:EtOAc ; 50:50) afforded the title compound.


Step 2 : 1-(3''-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}-1,1':2',1''-terphenyl-4-yl)ethanone.

Prepared according to the general procedure **Coupling-1** using bromide from step 1 and 4-acetylbenzeneboronic acid as starting material. Flash chromatography (Hex:EtOAc ; 50:50) afforded the title compound.

<sup>1</sup>H NMR (500 MHz, Benzene-*d*<sub>6</sub>): δ 8.81 (d, 1H), 7.93 (d, 1H); 7.85 (s, 1H), 7.80 (m, 3H), 7.75 (s, 1H), 7.61 (m, 2H), 7.38 (m, 7H), 6.87 (dd, 1H), 2.35 (s, 3H), 2.05 (s, 3H), 1.65 (s, 6H).  
+ESI (M+1) 520.2

#### EXAMPLE 99

2-(3''-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}-1,1':2',1''-terphenyl-4-yl)propan-2-ol



Step 1: 1-(2'-bromo-1,1'-biphenyl-4-yl)ethanone

MC054PV

Prepared according to the general procedure **Coupling-1** using 4-acetylbenzeneboronic acid and 1,2-dibromobenzene as starting material. Flash chromatography (Hex:EtOAc ; 95:5) afforded the title compound.

5 **Step 2: 2-(2'-bromo-1,1'-biphenyl-4-yl)propan-2-ol**

Prepared according to the general procedure **Cer-1** using methyl magnesium bromide and ketone from step 1 as starting material. Flash chromatography (Hex:EtOAc ; 90:10) afforded the title compound.

10

**Step 3: 2-(3''-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}-1,1':2',1''-terphenyl-4-yl)propan-2-ol**


15 Prepared according to the general procedure **Coupling-1** using **Quinoline\_4** and bromide from step 2 as starting material. Flash chromatography (Hex:EtOAc ; 50:50) afforded the title compound.

<sup>1</sup>H NMR (500 MHz, acetone-*d*<sub>6</sub>): δ 8.94 (dd, 1H); 8.45 (dd, 1H), 8.27 (d, 1H), 8.01 (d, 1H), 7.68 (d, 1H), 7.62 (br s, 1H), 7.55 (m, 2H), 7.48 (m, 3H), 7.42 (d, 2H), 7.38 (t, 1H), 7.21 (m, 3H).

20 +ESI (M+1) 536.8

**EXAMPLE 100**

25 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-{3-[5-(3-methyl-1,2,4-oxadiazol-5-yl)-1-oxidopyridin-3-yl]phenyl}quinoline



Prepared according to the general procedure **Coupling-1** using **Quinoline\_4** and **Pyr\_11** as starting material. Flash chromatography (EtOAc:EtOH ; 90:10) and crystallisation from EtOAc/ether afforded the title compound.

5

The following compounds (**Table 3**) were prepared according to the general procedure **Coupling\_1** using **Quinoline 4** and intermediates described previously or commercial products (Commercial). Indicated is their respective (M + 1)<sup>+</sup> value obtained from a low resolution mass spectrometer under electron-spray ionization conditions.

10

**Table 3:**

EX.	intermediate	Chemical name	ESI-LRMS (M + 1) <sup>+</sup>
101	Pyr_1	6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{5-[4-(methylsulfonyl)phenyl]-1-oxidopyridin-3-yl}phenyl)quinoline	573.1
102	Pyr_2	6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{5-[3-(methylsulfonyl)phenyl]-1-oxidopyridin-3-yl}phenyl)quinoline	573.6
103	Pyr_3	6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{5-[2-(methylsulfonyl)phenyl]-1-oxidopyridin-3-yl}phenyl)quinoline	573.2
104	Pyr_4	6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{2-[4-(methylsulfonyl)phenyl]pyridin-3-yl}phenyl)quinoline	557.1
105	Pyr_5	6-[1-methyl-1-(methylsulfonyl)ethyl]-8-[3-(1-oxido-5-phenylpyridin-3-yl)phenyl]quinoline	495.1
106	Pyr_6	8-{3-[5-(3,5-dichlorophenyl)-1-oxidopyridin-3-yl]phenyl}-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline	563.1/565.1
107	Pyr_7	8-{3-[5-(3,4-dimethoxyphenyl)-1-oxidopyridin-3-yl]phenyl}-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline	555.1
108	Pyr_8	6-[1-methyl-1-(methylsulfonyl)ethyl]-8-{3-[5-(5-methyl-1,2,4-oxadiazol-3-yl)-1-oxidopyridin-3-yl]phenyl}quinoline	501.1


109	Pyr_9	6-[1-methyl-1-(methylsulfonyl)ethyl]-8-{3-[5-(5-methyl-1,3,4-oxadiazol-2-yl)-1-oxidopyridin-3-yl]phenyl}quinoline	501.1
110	Pyr_10	8-{3-[6-(benzyloxy)-5-(3-methyl-1,2,4-oxadiazol-5-yl)pyridin-3-yl]phenyl}-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline	591.2
111	Pyr_12	6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{6-[4-(methylsulfonyl)phenyl]-1-oxidopyridin-3-yl}phenyl)quinoline	573.3
112	Pyr_13	6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{5-[4-(methylsulfonyl)phenyl]-1-oxidopyridin-2-yl}phenyl)quinoline	573.3
113	Pyr_14	6-[1-methyl-1-(methylsulfonyl)ethyl]-8-{3-[3-[4-(methylsulfonyl)phenyl]-5-(trifluoromethyl)pyridin-2-yl]phenyl}quinoline	625.3
114	Pyr_15	1-(4-chlorophenyl)-3-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)pyridin-2(1H)-one	529.1
115	Pyr_16	6-[1-methyl-1-(methylsulfonyl)ethyl]-8-{3-[5-(2-methyl-2H-tetraazol-5-yl)-1-oxidopyridin-3-yl]phenyl}quinoline	501.1
116	Het_1	N-isopropyl-5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-4-[4-(methylsulfonyl)phenyl]thiophene-2-carboxamide	647.4
117	Commercial	6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(1,1':2',1''-terphenyl-3-yl)quinoline	478.1
118	Phe_1	6-[1-methyl-1-(methylsulfonyl)ethyl]-8-[4''-(methylsulfonyl)-1,1':2',1''-terphenyl-3-yl]quinoline	556.2
119	Phe_2	6-[1-methyl-1-(methylsulfonyl)ethyl]-8-[2'-(5-methyl-1,3,4-oxadiazol-2-yl)-1,1'-biphenyl-3-yl]quinoline	484.2
120	Phe_3	methyl 3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}-4''-(methylsulfonyl)-1,1':2',1''-terphenyl-4'-carboxylate	
121	Commercial	6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(1,1':4',1''-terphenyl-3-yl)quinoline	478.2
122	Commercial	6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(1,1':3',1''-terphenyl-3-yl)quinoline	478.2

MC054PV

123	Phe_4	2-[5-(3'-(6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl)-1,1'-biphenyl-2-yl)-1,3-thiazol-2-yl]propan-2-ol	542.0
124	Phe_5	6-[1-methyl-1-(methylsulfonyl)ethyl]-8-[3'-(1-oxidopyridin-4-yl)-1,1'-biphenyl-3-yl]quinoline	495.1

### EXAMPLE 125

5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-3-(3-methyl-1,2,4-oxadiazol-5-yl)pyridin-2(1H)-one



5


A solution of **EXAMPLE 110** (1.0 eq) in  $\text{CH}_2\text{Cl}_2$  /TFA (2:1, 0.06M) was stirred at r.t. for 2 h, volatiles evaporated, diluted with EtOAc, washed with a solution of  $\text{Na}_2\text{CO}_3$  and brine. The organic extracts were dried over  $\text{MgSO}_4$ , filtered and concentrated. Stirring in EtOAc and isolation by filtration provided the title compound.

10

$^1\text{H}$  NMR (500 MHz,  $\text{DMSO}-d_6$ ):  $\delta$  8.92 (dd, 1H), 8.62 (d, 1H), 8.51 (dd, 1H), 8.27 (d, 1H), 8.17 (s, 1H), 8.03 (d, 1H), 7.86 (s, 1H), 7.68 (d, 1H), 7.62-7.55 (m, 3H), 2.80 (s, 3H), 2.37 (s, 3H), 1.92 (s, 6H).

## EXAMPLE 126

6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{6-[4-(methylsulfonyl)phenyl]imidazo[2,1-  
b][1,3]thiazol-5-yl}phenyl)quinoline



5

To a solution of the aldehydriketone from **EXAMPLE 1**, step 4 (1.0 eq) in DMF (0.1M) was added 2-aminothiazole (1.6 eq). The mixture was heated at 120°C for 1h, cooled, diluted with water, saturated NaHCO<sub>3</sub> solution and EtOAc. The organic extracts were washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated. The residue was purified by flash chromatography (Toluene:EtOAc, 1:3) to afford the title compound.

10


<sup>1</sup>H NMR (500 MHz, acetone-*d*<sub>6</sub>): δ 9.02 (dd, 1H), 8.50 (dd, 1H), 8.32 (d, 1H), 8.26 (d, 1H), 8.08-8.05 (m, 4H), 7.92 (d, 2H), 7.85 (d, 1H), 7.69 (t, 1H), 7.62-7.59 (m, 2H), 7.36 (d, 1H), 3.15 (s, 3H), 2.74 (s, 3H), 2.01 (s, 6H).

15

MC054PV

## EXAMPLE 127

6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{2-[4-(methylsulfonyl)phenyl]imidazo[1,2-  
a]pyridin-3-yl}phenyl)quinoline



5

Prepared according to the procedure described for **EXAMPLE 126** and using 2-aminopyridine as starting material. Flash chromatography (Toluene:EtOAc, 1:3) afforded the title compound.


<sup>1</sup>H NMR (500 MHz, acetone-*d*<sub>6</sub>): δ 9.00 (dd, 1H), 8.50 (d, 1H), 8.48 (dd, 1H), 8.32 (d, 1H), 8.27 (d, 1H), 8.14 (d, 2H), 8.07 (t, 1H), 7.94-7.92 (m, 3H), 7.76 (t, 1H), 7.67 (d, 1H), 7.62-7.59 (m, 2H), 7.39 (dd, 1H), 7.03 (dt, 1H), 3.16 (s, 3H), 2.74 (s, 3H), 2.01 (s, 6H).

10

15

## EXAMPLE 128

[3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}-4''-(methylthio)-1,1':2',1''-terphenyl-4'-yl]methanol



MC054PV

Step 1: 2-bromo-3'-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}-1,1'-biphenyl-4-carbaldehyde

Prepared according to the general procedure **Coupling\_1** using **Quinoline\_4** and 3,4-dibromobenzaldehyde as starting materials. Flash chromatography (hexane:EtOAc, 9:1 to 1:1) afforded the title compound as foam.

Step 2: 3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}-4''-(methylthio)-1,1':2',1''-terphenyl-4'-carbaldehyde

Prepared according to the general procedure **Coupling\_1** using aldehyde from step 1 and 4-methylthiobenzeneboronic acid as starting materials. Flash chromatography (hexane:EtOAc, 3:2 to 1:1) afforded the title compound.

Step 3: [3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}-4''-(methylthio)-1,1':2',1''-terphenyl-4'-yl]methanol


To a solution of aldehyde from step 2 in EtOH: THF (4:1, 0.05M) at 0°C was added NaBH<sub>4</sub> (1 eq). The reaction mixture was stirred 1 h at 0°C, quenched with a solution of NH<sub>4</sub>Cl and diluted with ether. The organic extracts were washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated. The residue was purified by flash chromatography (hexane:EtOAc, 1:1 to 1:9) to afford the title compound.

<sup>1</sup>H NMR (500 MHz, acetone-*d*<sub>6</sub>): δ 8.91 (dd, 1H), 8.43 (d, 1H), 8.25 (d, 1H), 7.88 (d, 1H), 7.67 (d, 1H), 7.56 (dd, 1H), 7.55-7.46 (m, 4H), 7.42 (t, 1H), 7.29-7.18 (m, 5H), 4.76 (d, 2H), 4.33 (t, 1H), 2.70 (s, 3H), 2.474 (s, 3H), 1.97 (s, 6H).



## EXAMPLE 129

[3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}-4''-(methylsulfonyl)-1,1':2',1''-terphenyl-4'-yl]methanol




Prepared according to the general procedure Oxid-1 using Example 128 as starting material. The title compound was isolated as white foam.

<sup>1</sup>H NMR (500 MHz, acetone-*d*<sub>6</sub>): δ 8.91 (dd, 1H), 8.44 (d, 1H), 8.26 (d, 1H), 7.98 (d, 1H), 7.85 (d, 2H), 7.68 (d, 1H), 7.58-7.52 (m, 7H), 7.41 (t, 1H), 7.25 (d, 1H), 4.79 (d, 2H), 4.4 (t, 1H), 3.03 (s, 3H), 2.71 (s, 3H), 1.97 (s, 6H).

## EXAMPLE 130

2-[3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}-4''-(methylsulfonyl)-1,1':2',1''-terphenyl-4'-yl]propan-2-ol



Prepared according to the general procedure Cer-1 using Example 120 as starting material. The residue was purified by flash chromatography (hexane:EtOAc, 1:1 to 3:7) to afford the title compound.


MC054PV

<sup>1</sup>H NMR (500 MHz, acetone-*d*<sub>6</sub>): δ 8.91 (dd, 1H), 8.42 (dd, 1H), 8.25 (d, 1H), 7.98 (d, 1H), 7.85 (d, 2H), 7.71-7.67 (m, 3H), 7.57-7.53 (m, 4H), 7.40 (t, 1H), 7.23 (d, 1H), 4.24 (s, OH), 3.03 (s, 3H), 2.71 (s, 3H), 1.97 (s, 6H), 1.63 (s, 6H).

5

## EXAMPLE 131

3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}-4''-(methylsulfonyl)-1,1':2',1''-terphenyl-4'-carboxylic acid



10

To a solution of **Example 120** in THF:MeOH (1:1, 0.05M) was added LiOH (2N, 5 eq). The reaction mixture was stirred at rt for 24h, acidified to pH 2 with 1N HCl and diluted with EtOAc. The organic extracts were washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated. The residue was sonicated in hexane/EtOAc and title product isolated by filtration as a white powder.

15


<sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD): δ 8.86 (dd, 1H), 8.44 (d, 1H), 8.22 (d, 1H), 8.16 (dd, 1H), 8.12 (d, 1H), 7.87 (d, 2H), 7.84 (d, 1H), 7.69 (d, 1H), 7.61-7.54 (m, 4H), 7.46 (t, 1H), 7.43 (s, 1H), 7.32 (d, 1H), 4.79 (d, 2H), 4.4 (t, 1H), 3.04 (s, 3H), 2.74 (s, 3H), 1.98 (s, 6H).

20

MC054PV

## EXAMPLE 132

6-[1-methyl-1-(methanesulfonyl)ethyl]-8-(3-{2-[4-(methanesulfonyl)phenyl]-1-oxidopyridin-3-yl}phenyl)quinoline



5


Prepared according to the general procedure Oxid-2 using Example 104 as starting material. The residue was sonicated in hexane/EtOAc and title product isolated by filtration as a white powder.

<sup>1</sup>H NMR (500 MHz, acetone-*d*<sub>6</sub>): δ 8.92 (dd, 1H), 8.42 (dd, 1H), 8.35 (d, 1H), 8.25 (d, 1H), 7.95 (d, 1H), 7.85 (d, 2H), 7.65 (d, 2H), 7.57-7.48 (m, 5H), 7.36 (t, 1H), 7.21 (d, 1H), 3.02 (s, 3H), 2.70 (s, 3H), 1.96 (s, 6H).

10

## EXAMPLE 133

8-{3-[3-(4-fluorophenyl)-1,2,4-oxadiazol-5-yl]phenyl}-6-[1-methyl-1-(methanesulfonyl)ethyl]quinoline



20

Step 1: methyl 3-{6-[1-methyl-1-(methanesulfonyl)ethyl]quinolin-8-yl}benzoate

To a solution of Quinoline\_2 in CH<sub>2</sub>Cl<sub>2</sub>:MeOH (1:2, 0.03M) was added NaCN (1.8 eq), AcOH (1 eq) followed by MnO<sub>2</sub> (8 eq). The reaction mixture was stirred at rt for

MC054PV

24h, filtered on Celite and diluted with water. The mixture was extracted with  $\text{CH}_2\text{Cl}_2$  (3X). The organic extracts were washed with brine, dried over  $\text{MgSO}_4$ , filtered and concentrated. The residue was purified by flash chromatography (hexane:EtOAc, 1:1) to afford the title compound.

5

Step 2: 3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}benzoic acid

To a solution of ester from step 1 in THF (0.08M) was added LiOH (2N, 5 eq). The reaction mixture was stirred at rt for 5h, acidified with AcOH, diluted with water and extracted with  $\text{CH}_2\text{Cl}_2$ . The organic extracts were washed with brine, dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated. The title compound was isolated by filtration from Hexane/ether as a solid.

10

Step 3: 8-{3-[3-(4-fluorophenyl)-1,2,4-oxadiazol-5-yl]phenyl}-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline

15

To a solution of acid from step 2 in DMF (0.05M) was added CDI (2 eq) and 4-fluoro-*N*'-hydroxybenzenecarboximidamide (2 eq). The reaction mixture was stirred at rt for 15 min and 18h at 120°C. The reaction mixture cooled to rt, diluted with water and extracted with EtOAc. The organic extracts were washed with brine, dried over  $\text{MgSO}_4$ , filtered and concentrated. The residue was purified by flash chromatography (hexane:EtOAc, 20:80) to afford the title compound.

20


$^1\text{H}$  NMR (500 MHz, acetone- $d_6$ ):  $\delta$  8.98 (dd, 1H), 8.64 (s, 1H), 8.55 (dd, 1H), 8.39 (d, 1H), 8.31 (d, 1H), 8.28 (d, 1H), 8.25 (dd, 2H), 8.08 (d, 1H), 7.82 (t, 1H), 7.63 (dd, 1H), 7.39 (t, 2H), 2.79 (s, 3H), 2.05 (s, 6H). +ESI (M+1) 488.2.

25

## EXAMPLE 134

6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{4-[4-(methylthio)phenyl]pyridin-3-yl}phenyl)quinoline

---



5

Step 1: 4-chloro-3-(tributylstannyl)pyridine

To a solution of LDA (1.1 eq) in THF (0.05M) at  $-78^{\circ}\text{C}$  was added 4-chloropyridine (1 eq). After 1.5h tributyltin chloride (1.5 eq) was added and the reaction mixture was warmed at rt over 3 h. The reaction mixture was diluted with water and extracted with EtOAc. The organic extracts were washed with brine, dried over  $\text{MgSO}_4$ , filtered and concentrated. The residue was purified by flash chromatography (hexane:EtOAc, 95:5) to afford the title compound.

10

+ESI (M+1) 525.3

15

Step 2: 8-[3-(4-chloropyridin-3-yl)phenyl]-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline

Prepared according to the general procedure **Coupling\_3** using **Quinoline\_3** and the Aryltin from step 1 as starting materials. Flash chromatography ( $\text{CH}_2\text{Cl}_2$ :MeOH, 99:1) afforded the title compound.

20

Step 3: 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{4-[4-(methylthio)phenyl]pyridin-3-yl}phenyl)quinoline

25

MC054PV


Prepared according to the general procedure **Coupling\_1** using 4-methylthiobenzeneboronic acid and the chloropyridine from step 2 as starting materials. Flash chromatography afforded the title compound.

- 5 <sup>1</sup>H NMR (500 MHz, acetone-*d*<sub>6</sub>): δ 8.91 (dd, 1H), 8.70 (s, 1H), 8.63 (d, 1H), 8.44 (dd, 1H), 8.27 (d, 1H), 7.92 (d, 1H), 7.76 (d, 1H), 7.58 (d, 1H), 7.56 (dd, 1H), 7.49 (t, 1H), 7.45 (d, 1H), 7.34 (d, 1H), 7.28 (d, 2H), 7.23 (d, 2H), 2.71 (s, 3H), 2.50 (s, 3H), 1.98 (s, 6H).  
+ESI (M+1) 525.3

10

#### EXAMPLE 135

3''-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}-1,1':2',1''-terphenyl-4-carboxylic acid



- 15 Step 1: 8-(2'-bromo-1,1'-biphenyl-3-yl)-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline

Prepared according to the general procedure **Coupling\_1** using **Quinoline\_4** and 1,2-dibromobenzene as starting materials. Flash chromatography (hexane:EtOAc, 1:1)

- 20 afforded the title compound.

Step 2: 3''-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}-1,1':2',1''-terphenyl-4-carboxylic acid


MC054PV

Prepared according to the general procedure **Coupling\_1** using Arylbromide from step 1 and 4-carboxybenzeneboronic acid as starting materials. Flash chromatography ( $\text{CH}_2\text{Cl}_2:\text{MeOH}$ , 9:1) afforded the title compound.

$^1\text{H}$  NMR (500 MHz, acetone- $d_6$ ):  $\delta$  8.90 (dd, 1H), 8.43 (dd, 1H), 8.24 (d, 2H), 8.07 (d, 2H), 7.96 (d, 1H), 7.83 (d, 1H), 7.68 (d, 1H), 7.62 (d, 1H), 7.52 (m, 4H), 7.41 (d, 2H), 7.28 (d, 1H), 2.68 (s, 3H), 1.95 (s, 6H). -ESI (M-1) 520.4

**EXAMPLE 136**

2-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-3-[4-(methylsulfonyl)phenyl]quinoxaline




A solution of 2-bromo-2-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-1-[4-(methylsulfonyl)phenyl]ethanone from **EXAMPLE 1**, step 4 in DMF (0.05M) and 1,2-diaminobenzene (2 eq) was stirred at 130°C for 18h. The reaction mixture was cooled to rt, diluted with water and extracted with EtOAc. The organic extracts were washed with brine, dried over  $\text{MgSO}_4$ , filtered and concentrated. The residue was purified by flash chromatography (hexane:EtOAc, 1:1) to afford the title compound.

$^1\text{H}$  NMR (500 MHz, acetone- $d_6$ ):  $\delta$  8.89 (dd, 1H), 8.43 (dd, 1H), 8.29 (d, 1H), 8.21 (m, 2H), 8.05 (d, 1H), 8.00 (d, 2H), 7.96 (d, 2H), 7.95 (m, 3H), 7.85 (d, 1H), 7.70 (d, 1H), 7.58 (m, 2H), 3.10 (s, 3H), 2.72 (s, 3H), 1.99 (s, 6H).

MC054PV

## EXAMPLE 137

6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{3-[4-(methylsulfonyl)phenyl]pyrazin-2-yl}phenyl)quinoline



5

Step 1: 8-[3-(3-chloropyrazin-2-yl)phenyl]-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline

10

Prepared according to the general procedure **Coupling\_1** using **Quinoline\_4** and 1,2-dichloropyrazine as starting materials. Flash chromatography (hexane:EtOAc, 3:7) afforded the title compound.

15

Step 2: 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{3-[4-(methylthio)phenyl]pyrazin-2-yl}phenyl)quinoline

Prepared according to the general procedure **Coupling\_1** using 4-methylthiobenzeneboronic acid and the chloropyrazine from step 1 as starting materials. Flash chromatography (hexane:EtOAc, 2:8) afforded the title compound.

20

Step 3: 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{3-[4-(methylsulfonyl)phenyl]pyrazin-2-yl}phenyl)quinoline

25

Prepared according to the general procedure **Oxid-1** using thioether from step 2 as starting material. Flash chromatography (hexane:EtOAc, 2:8) afforded the title compound.



<sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>): δ 8.86 (dd, 1H), 8.78 (d, 1H), 8.75 (d, 1H), 8.45 (dd, 1H), 8.27 (d, 1H), 8.02 (d, 1H), 7.94 (d, 2H), 7.85 (d, 2H), 7.83 (m, 2H), 7.60 (m, 1H), 7.39 (m, 1H), 7.25 (t, 1H), 3.09 (s, 3H), 2.71 (s, 3H), 1.98 (s, 6H). +ESI (M+1) 558.1

5

The following compounds (Table 4) were prepared according to the procedure described previously. Indicated is their respective (M + 1)<sup>+</sup> value obtained from a low resolution mass spectrometer under electron-spray ionization conditions.

10

Table 4:


EX.	Chemical name	ESI-LRMS (M + 1) <sup>+</sup>
138	6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{4-[4-(methylsulfonyl)phenyl]pyridin-3-yl}phenyl)quinoline	557.1
139	6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{4-[4-(methylsulfonyl)phenyl]-1-oxidopyridin-3-yl}phenyl)quinoline	573.3
140	6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{4-[3-(methylthio)phenyl]pyridin-3-yl}phenyl)quinoline	525.0
141	8-[4',5'-difluoro-4''-(methylthio)-1,1':2',1''-terphenyl-3-yl]-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline	560.1
142	8-[4',5'-difluoro-4''-(methylsulfonyl)-1,1':2',1''-terphenyl-3-yl]-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline	591.8
143	8-(4''-fluoro-1,1':2',1''-terphenyl-3-yl)-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline	496.1
144	6,7-dichloro-2-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-3-[4-(methylsulfonyl)phenyl]quinoxaline	676.1
145	2-(4-chlorophenyl)-3-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)quinoxaline	564.1
146	2-{4-[3-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)quinoxalin-2-yl]phenyl}propan-2-ol	588.0

MC054PV

147	2-[3,4-bis(difluoromethoxy)phenyl]-3-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)quinoxaline	662.0
148	4-[3-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)quinoxalin-2-yl]benzoic acid	574.3
149	<i>N</i> -cyclopropyl-4-[3-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)quinoxalin-2-yl]benzamide	613.4
150	2-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-3-(4-methylphenyl)quinoxaline	544.2
151	2-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-3-phenylquinoxaline	530.2
152	2-(4-fluorophenyl)-3-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)quinoxaline	548.8
153	2-{4-[3-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)pyrazin-2-yl]phenyl}propan-2-ol	536.8
154	6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{3-[4-(methylthio)phenyl]pyrazin-2-yl}phenyl)quinoline	526.3
155	8-{3-[3-(4-fluorophenyl)pyrazin-2-yl]phenyl}-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline	498.2
156	8-(3-{2-(2-ethylpyridin-4-yl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-5-yl}phenyl)-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline	668.3
157	2-(4-{5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-2-yl}phenyl)propan-2-ol	697.5

## EXAMPLE 158

5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-6-[4-(methylsulfonyl)phenyl]pyridin-2(1H)-one




5 To a suspension of ketone from **EXAMPLE 1**, step 3 in MeOH (0.15M) at 0°C was condensed NH<sub>3</sub> (30% volume) and added methyl propiolate (6 eq). The reaction mixture was heated in a pressure tube at 150°C for 5h, cooled to rt and concentrated to dryness. The residue was purified by flash chromatography (EtOH:EtOAc, 12:88) to afforded the title compound.

10 <sup>1</sup>H NMR (500 MHz, acetone-*d*<sub>6</sub>): δ 11.15 (br s, 1H), 8.87 (dd, 1H), 8.41 (dd, 1H), 8.23 (d, 1H), 7.95 (d, 1H), 7.88 (dd, 2H), 7.72 (d, 1H), 7.69 (dd, 2H), 7.63 (dt, 2H), 7.53 (m, 2H), 7.38 (t, 1H), 7.18 (dt, 1H), 6.59 (d, 1H), 3.04 (s, 3H), 2.69 (s, 3H), 1.94 (s, 6H).

15

## EXAMPLE 159

1-methyl-5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-6-[4-(methylsulfonyl)phenyl]pyridin-2(1H)-one



5

To a solution of **EXAMPLE 158** in DMF (0.05M) at 0°C was added NaH (1.1 eq) followed by MeI (6 eq). The reaction mixture was stirred at rt for 1h, diluted with NH<sub>4</sub>Cl solution and extracted with EtOAc. The organic extracts were washed with brine, dried over MgSO<sub>4</sub>, filtered and concentrated. The residue was purified by flash chromatography (hexane:EtOAc, 70:30, then EtOH:EtOAc, 12:88) to afford the title compound and **EXAMPLE 160**.


10

<sup>1</sup>H NMR (500 MHz, acetone-*d*<sub>6</sub>): δ 8.94 (dd, 1H), 8.45 (dd, 1H), 8.26 (d, 1H), 7.97 (m, 3H), 7.74 (d, 2H), 7.59 (m, 3H), 7.46 (t, 1H), 7.30 (t, 1H), 7.12 (d, 1H), 6.60 (d, 1H), 3.26 (s, 3H), 3.00 (s, 3H), 2.74 (s, 3H), 2.00 (s, 6H).

15

## EXAMPLE 160

8-(3-{6-methoxy-2-[4-(methylsulfonyl)phenyl]pyridin-3-yl}phenyl)-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline



5


Prepared according to the procedure described for **EXAMPLE 159**.

<sup>1</sup>H NMR (500 MHz, acetone-*d*<sub>6</sub>): δ 8.89 (dd, 1H), 8.44 (dd, 1H); 8.26 (d, 1H), 8.00 (d, 1H), 7.90 (d, 1H), 7.87 (d, 2H), 7.79 (d, 2H), 7.71 (dt, 1H), 7.61 (t, 1H), 7.56 (dd, 1H), 7.48 (t, 1H), 7.29 (dt, 1H), 6.96 (d, 1H), 4.03 (s, 3H), 3.10 (s, 3H), 2.71 (s, 3H), 1.97 (s, 6H).

10

## EXAMPLE 161

8-(3-{6-(difluoromethoxy)-2-[4-(methylsulfonyl)phenyl]pyridin-3-yl}phenyl)-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline



15


Prepared according to procedure described for **EXAMPLE 159** but using ethyl bromodifluoroacetate as electrophile. Flash chromatography (hexane:EtOAc, 70:30, then EtOH:EtOAc, 12:88) afforded the title compound.

<sup>1</sup>H NMR (500 MHz, acetone-*d*<sub>6</sub>): δ 8.89 (dd, 1H), 8.45 (dd, 1H), 8.27 (d, 1H), 8.13 (d, 1H), 8.01 (d, 1H), 7.89 (d, 2H), 7.87 (t, 1H), 7.80 (d, 2H), 7.76 (dt, 1H), 7.65 (t, 1H), 7.57 (dd, 1H), 7.51 (t, 1H), 7.34 (dt, 1H), 7.22 (d, 2H), 3.11 (s, 3H), 2.71 (s, 3H), 1.97 (s, 6H).

5

**EXAMPLE 162**

8-(3-{6-[(4-fluorobenzyl)oxy]-2-[4-(methylsulfonyl)phenyl]pyridin-3-yl}phenyl)-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline




10

Prepared according to procedure described for **EXAMPLE 159** but using 4-fluorobenzyl bromide as electrophile. Flash chromatography (hexane:EtOAc, 70:30, EtOH:EtOAc, 12:88, EtOH:EtOAc, 5:95) afforded the title compound and **EXAMPLE 162**.

15 <sup>1</sup>H NMR (500 MHz, acetone-*d*<sub>6</sub>): δ 8.89 (dd, 1H), 8.44 (dd, 1H), 8.26 (d, 1H), 8.01 (d, 1H), 7.92 (d, 1H), 7.88 (d, 2H), 7.78 (d, 2H), 7.72 (dt, 1H), 7.61 (m, 3H), 7.56 (dd, 1H), 7.47 (t, 1H), 7.30 (dt, 1H), 7.19 (t, 2H), 7.02 (d, 1H), 5.53 (s, 2H), 3.10 (s, 3H), 2.71 (s, 3H), 1.97 (s, 6H).

## EXAMPLE 163

1-(4-fluorobenzyl)-5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-  
6-[4-(methylsulfonyl)phenyl]pyridin-2(1H)-one



5


Prepared according to the procedure described for EXAMPLE 162.

<sup>1</sup>H NMR (500 MHz, acetone-*d*<sub>6</sub>): δ 8.93 (dd, 1H), 8.44 (dd, 1H), 8.25 (d, 1H), 7.92 (d, 1H), 7.82 (d, 2H), 7.68 (d, 1H), 7.57 (m, 2H), 7.47 (d, 2H), 7.43 (s, 1H), 7.29 (t, 1H), 7.14 (d, 2H), 6.98 (t, 2H), 6.92 (m, 2H), 6.73 (d, 1H), 5.18 (br s, 2H), 2.99 (s, 3H), 2.73 (s, 3H), 1.98 (s, 6H).

10

## EXAMPLE 164

5-(4-fluorophenyl)-6-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)pyridin-2(1H)-one



15

Step 1: 2-(4-fluorophenyl)-1-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)ethanone

MC054PV

Prepared according to the procedure described for **EXAMPLE 1**, step 1-2, using **Quinoline\_2** in step 1 and 4-fluorobenzaldehyde in step 2. Flash chromatography (Hexane:EtOAc, 40:60) afforded the title compound.


Step 2: 5-(4-fluorophenyl)-6-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)pyridin-2(1*H*)-one

Prepared according to the procedure described for **EXAMPLE 158**. Flash chromatography (EtOH:EtOAc, 12:88) afforded the title compound.

<sup>1</sup>H NMR (500 MHz, Acetone/Chloroform = 2:1):  $\delta$  8.89 (dd, 1H), 8.39 (dd, 1H), 8.22 (d, 1H), 7.88 (s, 1H), 7.79 (d, 2H), 7.62 (s, 1H), 7.57 (d, 1H), 7.54 (dd, 1H), 7.46 (t, 1H), 7.38 (d, 1H), 7.22 (m, 2H), 7.02 (t, 2H), 6.52 (d, 2H), 2.67 (s, 3H), 1.96 (s, 6H).

#### EXAMPLE 165

5-(4-fluorophenyl)-1-methyl-6-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)pyridin-2(1*H*)-one



20

Prepared according to the procedure described for **EXAMPLE 159** using **EXAMPLE 164** as starting material. The residue was purified by flash chromatography (hexane:EtOAc, 60:40, then EtOH:EtOAc, 8:92) to afford the title compound and **EXAMPLE 166**.




MC054PV

<sup>1</sup>H NMR (500 MHz, acetone-*d*<sub>6</sub>): δ 8.97 (dd, 1H), 8.47 (dd, 1H), 8.31 (d, 1H), 8.05 (d, 1H), 7.98 (s, 1H), 7.80 (s, 1H), 7.77 (d, 1H), 7.60 (dd, 1H), 7.50 (m, 2H), 7.31 (d, 1H), 7.26 (m, 2H), 6.98 (t, 2H), 6.54 (d, 2H), 3.38 (s, 3H), 2.73 (s, 3H), 2.01 (s, 3H), 1.99 (s, 3H).

5

## EXAMPLE 166

8-{3-[3-(4-fluorophenyl)-6-methoxypyridin-2-yl]phenyl}-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline



10


Prepared according to the procedure described for EXAMPLE 165.

<sup>1</sup>H NMR (500 MHz, acetone-*d*<sub>6</sub>): δ 8.90 (dd, 1H), 8.44 (dd, 1H), 8.27 (dd, 1H), 7.96 (dd, 1H), 7.80 (t, 1H), 7.75 (d, 1H), 7.72 (dt, 1H), 7.57 (dd, 1H), 7.48 (dt, 1H), 7.41 (t, 1H), 7.32 (m, 2H), 7.11 (t, 2H), 6.87 (d, 1H), 4.01 (s, 3H), 2.70 (s, 3H), 1.98 (s, 6H).

15

## EXAMPLE 167

*N*-cyclopropyl-8-(3-{2-(1-hydroxy-1-methylethyl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-5-yl}phenyl)quinoline-6-carboxamide



5

Step 1: 2-{5-(3-bromophenyl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-2-yl}propan-2-ol

10 Prepared according to the procedure described for **EXAMPLE 71** using 3-bromo benzaldehyde as precursor. Flash chromatography (Hexane:EtOAc, 50:50-20:80) afforded the title compound.

Step 2: 2-{5-[3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-phenyl]-4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-2-yl}propan-2-ol

15

Prepared according to the procedure described for **Quinoline\_4** using bromide from step 1. Flash chromatography (Hexane:EtOAc, 50:50) afforded the title compound.

20 Step 3: *N*-cyclopropyl-8-(3-{2-(1-hydroxy-1-methylethyl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-5-yl}phenyl)quinoline-6-carboxamide

Prepared according to the procedure **Coupling-1** using boronic ester from step 2 and **Quinoline\_5** as starting material. Flash chromatography (Hex:EtOAc ; 10:90) afforded the title compound.


MC054PV

<sup>1</sup>H NMR (500 MHz, acetone-*d*<sub>6</sub>): δ 8.94 (dd, 1H), 8.48 (br s, 1H), 8.45 (d, 1H), 8.22 (d, 1H), 8.16 (s, NH), 7.94 (m, 4H), 7.81 (m, 2H), 7.62 (t, 1H), 7.59 (d, 1H), 7.50 (d, 1H), 5.21 (s, OH), 3.19 (s, 3H), 3.00 (s, 1H), 1.71 (s, 6H), 0.80 (m, 2H), 0.68 (m, 2H).

5

## EXAMPLE 168

8-(3-(2-(1-hydroxy-1-methylethyl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-5-yl)phenyl)quinoline-6-carboxylic acid




10 Prepared according to the procedure **Coupling-1** using boronic ester from **EXAMPLE 167**, step 2 and **8-bromoquinoline-6-carboxylic acid** from **Quinoline\_5**, step 1 as starting material. Flash chromatography (CH<sub>2</sub>Cl<sub>2</sub>:MeOH; 90:10) afforded the title compound.

15 <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD): δ 9.35 (d, 1H), 9.17 (dd, 1H), 9.03 (br s, 1H), 8.43 (br s, 1H), 8.18 (m, 1H), 7.93 (d, 2H), 7.87 (d, 2H), 7.69 (m, 4H), 3.40 (s, OH), 3.17 (s, 3H), 1.71 (s, 6H). +ESI, Q1 (M+1) 545.2

MC054PV

## EXAMPLE 169

2-[4-[4-(methylsulfonyl)phenyl]-5-(3-{6-[1-methyl-1-(1*H*-tetrazol-5-yl)ethyl]quinolin-8-yl}phenyl)-1,3-thiazol-2-yl]propan-2-ol



5

**Step 1: 8-bromo-6-[1-methyl-1-(1*H*-tetrazol-5-yl)ethyl]quinoline**

To a solution of Quinoline\_6 in Xylenes (0.2M) was added tributyltin chloride (5 eq) and sodium azide (5 eq). The reaction mixture was stirred at 135°C for 2 days.

- 10 The reaction mixture was diluted with NaOH 5N and the solid isolated by filtration washing with benzene/hexane. The material was suspended and stirred in MeOH/Et<sub>2</sub>O and the title product isolated by filtration.

15 **Step 2: 2-[4-[4-(methylsulfonyl)phenyl]-5-(3-{6-[1-methyl-1-(1*H*-tetrazol-5-yl)ethyl]quinolin-8-yl}phenyl)-1,3-thiazol-2-yl]propan-2-ol**


Prepared according to the procedure **Coupling-1** using boronic ester from EXAMPLE 167, step 2 and tetrazole from step 1 as starting material. Flash chromatography (CH<sub>2</sub>Cl<sub>2</sub>:MeOH:NH<sub>4</sub>OH; 90:10:5) afforded the title compound.

- 20 <sup>1</sup>H NMR (500 MHz, acetone-*d*<sub>6</sub>): δ 8.82 (dd, 1H), 8.33 (dd, 1H), 7.93 (m, 4H), 7.87 (d, 1H), 7.76 (br s, 1H), 7.72 (d, 1H), 7.69 (d, 1H), 7.50 (m, 2H), 7.42 (d, 1H), 5.51 (br s, NH), 3.32 (s, OH), 3.15 (s, 3H), 1.98 (s, 6H), 1.71 (s, 6H). +ESI, Q1 (M+1) 611.3

MC054PV

## EXAMPLE 170

1,1,1,3,3,3-hexafluoro-2-[8-(3-{2-(1-hydroxy-1-methylethyl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-5-yl}phenyl)quinolin-6-yl]propan-2-ol



5

Step 1: 3-bromo-4-(hexafluoro-2-hydroxyisopropyl)aniline

To a solution of 4-(hexafluoro-2-hydroxyisopropyl)aniline in DMF (0.08M) was added NBS (1.1 eq). The reaction mixture was stirred at rt for 1h, diluted with water and extracted with EtOAc. The organic extracts were washed with brine, dried over MgSO<sub>4</sub>, filtered and concentrated to afford the title compound.

15

Step 2: 2-(8-bromoquinolin-6-yl)-1,1,1,3,3,3-hexafluoropropan-2-ol

Prepared according to the procedure described for Quinoline-5, step 1 using aniline from step 1 as starting material. Flash chromatography (Hex:EtOAc ; 80:20 to 50:50) afforded the title compound.

20

Step 3: 1,1,1,3,3,3-hexafluoro-2-[8-(3-{2-(1-hydroxy-1-methylethyl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-5-yl}phenyl)quinolin-6-yl]propan-2-ol

MC054PV

Prepared according to the procedure **Coupling-1** using boronic ester from **EXAMPLE 167**, step 2 and **quinoline** from step 1 as starting material. Flash chromatography (Hex:EtOAc ; 80:20 to 50:50) afforded the title compound.


<sup>1</sup>H NMR (500 MHz, acetone-*d*<sub>6</sub>): δ 8.98 (d, 1H), 8.60 (d, 1H), 8.46 (s, 1H), 8.13 (s, 1H), 7.94 (m, 4H), 7.82 (m, 2H), 7.66 (dd, 1H), 7.61 (t, 1H), 7.51 (d, 1H), 5.21 (s, OH), 3.12 (s, 3H), 2.87 (s, OH), 2.05 (s, 3H), 1.72 (s, 6H). +ESI, Q1 (M+1) 667.4

10

MC054PV

## WHAT IS CLAIMED IS:

1. A compound represented by Formula (I):



(I)

or a pharmaceutically acceptable salt, wherein  
S1, S2, and S3 are independently

1. H,
2. -OH,
3. halogen,
4. -C<sub>1</sub>-C<sub>6</sub>alkyl,
5. -O-C<sub>1</sub>-C<sub>6</sub>alkyl optionally substituted with 1, 2 or 3 halogens, or -CN;

R<sub>1</sub> is

1. -(C<sub>1</sub>-C<sub>6</sub>alkyl)-SO<sub>n</sub>-(C<sub>1</sub>-C<sub>6</sub>alkyl) group, optionally substituted with 1, 2 or 3 substituents; wherein each substituent is independently a halogen, -OH and -CN,
2. -C(O)-O-aryl,
3. -C(O)-NH-aryl,
4. -C(O)-NH-heterocycle or N-oxide thereof,
5. -C(O)-NH-C<sub>1</sub>-C<sub>6</sub>alkyl,
6. -C(O)-NH-cycloC<sub>3</sub>-C<sub>6</sub>alkyl,
7. -C<sub>1</sub>-C<sub>6</sub>alkyl, optionally substituted with 1 to 6 halogens and 1 hydroxy,
8. -COOH,
9. -C<sub>1</sub>-C<sub>6</sub>alkyl-COOH,
10. -O-C<sub>1</sub>-C<sub>6</sub>alkyl,

MC054PV

11. -cycloC<sub>3</sub>-C<sub>6</sub>alkyl,
12. -C<sub>3</sub>-C<sub>6</sub>alkyl-heterocycle,
13. aryl,
14. heterocycle,
15. carbonyl,
16. carbamoyl, or
17. -SO<sub>n</sub>-(C<sub>1</sub>-C<sub>6</sub>alkyl);

each n is independently 0, 1, or 2;

Ar<sub>1</sub> and Ar<sub>2</sub> are each independently an aryl or heterocycle or an N-oxide thereof;

R<sub>2</sub> is

1. Hydrogen,
2. aryl optionally substituted with 1, 2 or 3 substituents selected from halogen,
3. heterocycle optionally substituted with 1, 2 or 3 halogens,
4. -C<sub>1</sub>-C<sub>6</sub>alkyl optionally substituted with 1, 2 or 3 substituents selected from hydroxy and halogen,
5. -COOH,
6. 1, 2 or 3 halogens,
7. -SO<sub>n</sub>-(C<sub>1</sub>-C<sub>6</sub>alkyl),
8. -N(H)-S(O)<sub>n</sub>-C<sub>1</sub>-C<sub>6</sub>alkyl,
9. -O-C<sub>1</sub>-C<sub>6</sub>alkyl substituents each optionally substituted with 1, 2 or 3 halogens,
10. -C(O)-N(H)-C<sub>3</sub>-C<sub>6</sub>cycloalkyl, or
11. -C(O)-C<sub>1</sub>-C<sub>6</sub>alkyl;

R<sub>3</sub> is

1. Hydrogen,
2. -C<sub>1</sub>-C<sub>6</sub>alkyl optionally substituted with hydroxy, -S(O)<sub>n</sub>-C<sub>1</sub>-C<sub>6</sub>alkyl, heterocycle, or 1, 2, 3, 4, 5 or 6 halogens,
3. aryl or C<sub>6</sub>-C<sub>12</sub>cycloalkyl optionally substituted with phenyl, -C<sub>1</sub>-C<sub>6</sub>alkyl, -S(O)<sub>n</sub>-C<sub>1</sub>-C<sub>6</sub>alkyl, -C(O)-O-C<sub>1</sub>-C<sub>6</sub>alkyl, -COOH, hydroxy-C<sub>1</sub>-C<sub>6</sub>alkyl- or 1, 2 or 3 halogens,



4. heterocycle or optionally substituted with 1, 2 or 3 substituents independently selected from phenyl, halogen, C<sub>1</sub>-C<sub>6</sub>alkyl, hydroxyC<sub>1</sub>-C<sub>6</sub>alkyl, -COOH, -C(O)-O-C<sub>1</sub>-C<sub>6</sub>alkyl,
5. amino,
6. -C(O)-O-C<sub>1</sub>-C<sub>6</sub>alkyl,
7. -C<sub>1</sub>-C<sub>6</sub>alkyl-O-phenyl optionally substituted with 1, 2 or 3 halogens,
8. -C<sub>1</sub>-C<sub>6</sub>alkyl-phenyl optionally substituted with 1 or 2 substituents selected from hydroxy and halo,
9. -COOH,
10. Halogen,
11. -SO<sub>n</sub>-(C<sub>1</sub>-C<sub>6</sub>alkyl),
12. -N(H)-S(O)<sub>n</sub>-C<sub>1</sub>-C<sub>6</sub>alkyl optionally substituted with 1, 2 or 3 halogen,
13. -N(H)-C(O)-C<sub>1</sub>-C<sub>6</sub>alkyl,
14. -N(H)-heterocycle optionally substituted with 1, 2 or 3 halogens,
15. -N(H)-aryl optionally substituted with 1, 2 or 3 halogens,
16. -N(H)-C<sub>1</sub>-C<sub>6</sub>alkyl optionally substituted with 1, 2 or 3 halogens,
17. -C(O)-N(H)-C<sub>1</sub>-C<sub>6</sub>alkyl optionally substituted with 1, 2 or 3 halogens,
18. -C(O)-NH-C<sub>3</sub>-C<sub>6</sub>cycloalkyl,
19. -O-C<sub>1</sub>-C<sub>6</sub>alkyl optionally substituted with 1, 2 or 3 halogens or phenyl optionally substituted with 1, 2, or 3 halogen;
- R<sub>4</sub> is
1. H,
2. Halogen,
3. -CN
4. -C<sub>1</sub>-C<sub>6</sub>alkyl,
5. -O-C<sub>1</sub>-C<sub>6</sub>alkyl optionally substituted with 1, 2 or 3 halogens,
6. -C<sub>1</sub>-C<sub>6</sub>alkyl-phenyl with phenyl optionally substituted with halogen, or
7. Oxo.

2. A compound according to claim 1, or a pharmaceutically acceptable salt thereof, wherein

MC054PV

Ar<sub>1</sub> is pyridine or pyridinone or an N-oxide thereof.

3. A compound according to claim 2, or a pharmaceutically acceptable salt thereof, wherein

Ar<sub>2</sub> is phenyl, oxadiazole or thiadiazole.

4. A compound according to claim 3, or a pharmaceutically acceptable salt thereof, wherein

R<sub>1</sub> is -(C<sub>1</sub>-C<sub>6</sub>alkyl)-SO<sub>n</sub>-(C<sub>1</sub>-C<sub>6</sub>alkyl); and

R<sub>2</sub> is -SO<sub>n</sub>-C<sub>1</sub>-C<sub>6</sub>alkyl.

5. A compound according to claim 1, or a pharmaceutically acceptable salt thereof, wherein

Ar<sub>1</sub> is phenyl.

6. A compound according to claim 5, or a pharmaceutically acceptable salt thereof, wherein

Ar<sub>2</sub> is phenyl, oxadiazole, thiadiazole, pyridine or pyridinone or an N-oxide thereof.

7. A compound according to claim 6, or a pharmaceutically acceptable salt thereof, wherein

R<sub>1</sub> is -(C<sub>1</sub>-C<sub>6</sub>alkyl)-SO<sub>n</sub>-(C<sub>1</sub>-C<sub>6</sub>alkyl); and

R<sub>2</sub> is -SO<sub>n</sub>-C<sub>1</sub>-C<sub>6</sub>alkyl.

8. A compound according to claim 1, or a pharmaceutically acceptable salt thereof, wherein

Ar<sub>1</sub> is thiazole or oxazole.

9. The compound according to claim 8, or a pharmaceutically acceptable salt thereof, wherein

Ar<sub>2</sub> is phenyl, pyridine or pyridinone or an N-oxide thereof.

MC054PV


10. The compound according to claim 9, or a pharmaceutically acceptable salt thereof, wherein

R<sub>1</sub> is -(C<sub>1</sub>-C<sub>6</sub>alkyl)-SO<sub>n</sub>-(C<sub>1</sub>-C<sub>6</sub>alkyl); and

R<sub>2</sub> is -SO<sub>n</sub>-C<sub>1</sub>-C<sub>6</sub>alkyl.

5

11. The compound according to claim 1 of Formula Ia



10

Ia

or a pharmaceutically acceptable salt thereof, wherein:

Ar<sub>1</sub> is phenyl, pyridine, pyridinone, pyrimidyl, thiophene, thiazole, triazole, tetrazole, oxazole, thiaphendiazole, pyridindiazole, imidazothiazole or quinoxaline or an N-oxide thereof; and

15

Ar<sub>2</sub> is phenyl, pyridine, pyridinone, oxadiazole or thiadiazole or an N-oxide thereof.

12. A compound according to claim 11, or a pharmaceutically acceptable salt thereof, wherein:

20

R<sub>2</sub> is phenyl, -COOH, -C<sub>1</sub>-C<sub>6</sub>alkyl, -C<sub>1</sub>-C<sub>6</sub>alkoxy, mono or di-halo-C<sub>1</sub>-C<sub>6</sub>alkoxy, hydroxyC<sub>1</sub>-C<sub>6</sub>alkyl, or -SO<sub>n</sub>-(C<sub>1</sub>-C<sub>6</sub>alkyl) or 1, 2 or 3 halogens;

13. A compound according to claim 12, or a pharmaceutically acceptable salt thereof, wherein:

25

MC054PV

R<sub>3</sub> is Hydrogen, amino, biphenyl, N-(tert-butoxycarbonyl)-4-phenylpyrrolidin-3-yl, N-(tert-butoxycarbonyl)azetidin-3-yl, N-(tert-butoxycarbonyl)pyrrolidin-3-yl, 3-chloro-4-fluorophenyl, 4-chlorophenoxyethyl, 2-chlorophenyl, 4-chlorophenyl, ethoxycarbonyl, furan-2-yl, furan-3-yl, imidazol-2-yl, indan-1-yl, indan-2-yl, 1H-indol-2-yl, 1H-indol-3-yl, 1H-indol-4-yl, 1H-indol-5-yl, 1H-indol-6-yl, 1H-indol-7-yl, isoquinolin-1-yl, isoquinolin-4-yl, isoquinolin-5-yl, isoquinolin-8-yl, isoxazol-3-yl, 3-methoxycarbonylphenyl, 4-methoxycarbonylphenyl, methyl, 1-methyl-1Hpyrazol-3-yl, 1-methyl-1Hpyrazol-4-yl, 1-methyl-1Hpyrazol-5-yl, 2-methylphenyl, 3-methylphenyl, 4-methylphenyl, 2-methylpyridin-5-yl, methylsulfonylmethyl, 2-methylsulfonylphenyl, 3-methylsulfonylphenyl, 4-methylsulfonylphenyl, morpholin-4-ylmethyl, phenyl, pyrazinyl, 1Hpyrazol-3-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, 3-pyridinylmethyl, pyrimidin-2-yl, pyrimidin-4-yl, pyrimidin-5-yl, quinolin-4-yl, quinolin-5-yl, quinolin-8-yl, 5,6,7,8-tetrahydro-1,8-naphthyridin-2-yl, 5,6,7,8-tetrahydro-5Hbenzo[a][7]annulen-5-yl, 5,6,7,8-tetrahydro-5Hbenzo[a][7]annulen-6-yl, tetrahydrofuran-2-yl, 1,2,3,4-tetrahydronaphthalen-1-yl, 1,2,3,4-tetrahydronaphthalen-2-yl, 1,3-thiazol-2-yl, 1,3-thiazol-5-yl, thiophen-2-yl and thiophen-2-yl.

14. A compound according to claim 11, or a pharmaceutically acceptable salt thereof, wherein:

Ar<sub>1</sub> is thiazole;

Ar<sub>2</sub> is phenyl; and

R<sub>2</sub> is -SO<sub>2</sub>-C<sub>1</sub>-C<sub>6</sub>alkyl or halogen or C<sub>1</sub>-C<sub>6</sub>alkyl optionally substituted with hydroxy or 1-3 halogens.

15. A compound according to claim 14 or a pharmaceutically acceptable salt thereof, wherein:

R<sub>3</sub> is Hydrogen or -C<sub>1</sub>-C<sub>6</sub>alkyl optionally substituted with hydroxy, -S(O)<sub>n</sub>C<sub>1</sub>-C<sub>6</sub>alkyl, or 1-6 halogens.

16. A compound according to claim 11, or a pharmaceutically acceptable salt thereof, wherein

Ar<sub>1</sub> is pyridine or an N-oxide thereof;

Ar<sub>2</sub> is oxadiazole; and

R<sub>2</sub> is

1. -C<sub>1</sub>-C<sub>6</sub>alkyl optionally substituted with hydroxy, -S(O)<sub>n</sub>C<sub>1</sub>-C<sub>6</sub>alkyl, or 1-3 substituents halogens,
2. -N(H)-C(O)-C<sub>1</sub>-C<sub>6</sub>alkyl,
3. -COOH, or
4. -C(O)-NH-C<sub>3</sub>-C<sub>6</sub>cycloalkyl.

17. A compound according to claim 16, or a pharmaceutically acceptable salt thereof wherein:  
R<sub>3</sub> is hydrogen.

18. The compound according to claim 1, selected from the group consisting of:

8-(3-{2-(3-chlorophenyl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-5-yl}phenyl)-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline,  
6-[1-methyl-1-(methylsulfonyl)ethyl]-8-{3-[4-[4-(methylsulfonyl)phenyl]-2-(1-oxidopyridin-4-yl)-1,3-thiazol-5-yl]phenyl}quinoline,  
6-[1-methyl-1-(methylsulfonyl)ethyl]-8-{3-[4-[4-(methylsulfonyl)phenyl]-2-(1-oxidopyridin-3-yl)-1,3-thiazol-5-yl]phenyl}quinoline,  
2-(3-{5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-2-yl}phenyl)propan-2-ol,  
3-{5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-2-yl}benzoic acid,  
2-{5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-2-yl}propan-2-ol,  
6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{2-(3-methyl-1,2,4-oxadiazol-5-yl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-5-yl}phenyl)quinoline,  
6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-5-yl}phenyl)quinoline,  
*N*-cyclopropyl-5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazole-2-carboxamide,

MC054PV


- 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{2-(6-methyl-1-oxidopyridin-3-yl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-5-yl}phenyl)quinoline,  
 2-[4-(4-chlorophenyl)-5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-1,3-thiazol-2-yl]propan-2-ol,  
 5 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{2-methyl-4-[4-(methylsulfonyl)phenyl]-1,3-oxazol-5-yl}phenyl)quinoline,  
 2-{4-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-5-[4-(methylsulfonyl)phenyl]-1,3-thiazol-2-yl}propan-2-ol,  
 10 1,1,1-trifluoro-*N*-(5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-2-yl)methanesulfonamide,  
 2-[5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-4-pyridin-3-yl-1,3-thiazol-2-yl]propan-2-ol,  
 2-[5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-4-(1-oxidopyridin-3-yl)-1,3-thiazol-2-yl]propan-2-ol,  
 15 1-(4-chlorophenyl)-1-{4-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-5-[4-(methylsulfonyl)phenyl]-1,3-thiazol-2-yl}ethanol,  
 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{3-[4-(methylsulfonyl)phenyl]thien-2-yl}phenyl)quinoline,  
 1-(3''-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}-1,1':2',1''-terphenyl-4-yl)ethanone,  
 20 2-(3''-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}-1,1':2',1''-terphenyl-4-yl)propan-2-ol,  
 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-{3-[5-(3-methyl-1,2,4-oxadiazol-5-yl)-1-oxidopyridin-3-yl]phenyl}quinoline,  
 25 5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-3-(3-methyl-1,2,4-oxadiazol-5-yl)pyridin-2(1*H*)-one,  
 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{6-[4-(methylsulfonyl)phenyl]imidazo[2,1-*b*][1,3]thiazol-5-yl}phenyl)quinoline,  
 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{2-[4-(methylsulfonyl)phenyl]imidazo[1,2-*a*]pyridin-3-yl}phenyl)quinoline  
 30 [3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}-4''-(methylthio)-1,1':2',1''-terphenyl-4'-yl]methanol,  
 [3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}-4''-(methylsulfonyl)-1,1':2',1''-terphenyl-4'-yl]methanol,

MC054PV

- 2-[3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}-4''-(methylsulfonyl)-1,1':2',1''-terphenyl-4'-yl]propan-2-ol,  
 3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}-4''-(methylsulfonyl)-1,1':2',1''-terphenyl-4'-carboxylic acid,  
 5 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{2-[4-(methylsulfonyl)phenyl]-1-oxidopyridin-3-yl}phenyl)quinoline,  
 8-{3-[3-(4-fluorophenyl)-1,2,4-oxadiazol-5-yl]phenyl}-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline,  
 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{4-[4-(methylthio)phenyl]pyridin-3-yl}phenyl)quinoline,  
 10 3''-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}-1,1':2',1''-terphenyl-4-carboxylic acid,  
 2-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-3-[4-(methylsulfonyl)phenyl]quinoxaline,  
 15 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{3-[4-(methylsulfonyl)phenyl]pyrazin-2-yl}phenyl)quinoline,  
 5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-6-[4-(methylsulfonyl)phenyl]pyridin-2(1H)-one,  
 1-methyl-5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-6-[4-(methylsulfonyl)phenyl]pyridin-2(1H)-one,  
 20 8-(3-{6-methoxy-2-[4-(methylsulfonyl)phenyl]pyridin-3-yl}phenyl)-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline,  
 8-(3-{6-(difluoromethoxy)-2-[4-(methylsulfonyl)phenyl]pyridin-3-yl}phenyl)-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline,  
 25 8-(3-{6-[(4-fluorobenzyl)oxy]-2-[4-(methylsulfonyl)phenyl]pyridin-3-yl}phenyl)-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline,  
 1-(4-fluorobenzyl)-5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-6-[4-(methylsulfonyl)phenyl]pyridin-2(1H)-one,  
 5-(4-fluorophenyl)-6-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)pyridin-2(1H)-one,  
 30 5-(4-fluorophenyl)-1-methyl-6-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)pyridin-2(1H)-one,  
 8-{3-[3-(4-fluorophenyl)-6-methoxypyridin-2-yl]phenyl}-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline,  
 35 or a pharmaceutically acceptable salt thereof.

MC054PV

19. The compound according to claim 1, of formula Ia:



wherein

R is selected from the group consisting of
Amino
2-biphenyl
3-biphenyl
N-(tert-butoxycarbonyl)-4-phenylpyrrolidin-3-yl
N-(tert-butoxycarbonyl)azetidin-3-yl
N-(tert-butoxycarbonyl)pyrrolidin-3-yl
3-chloro-4-fluorophenyl
4-chlorophenoxymethyl
2-chlorophenyl
4-chlorophenyl
Ethoxycarbonyl
furan-2-yl
furan-3-yl
imidazol-2-yl
indan-1-yl
indan-2-yl
1H-indol-2-yl
1H-indol-3-yl
1H-indol-4-yl
1H-indol-5-yl
1H-indol-6-yl
1H-indol-7-yl
Isoquinolin-1-yl



MC054PV

Isoquinolin-4-yl
Isoquinolin-5-yl
Isoquinolin-8-yl
isoxazol-3-yl
3-methoxycarbonylphenyl
4-methoxycarbonylphenyl
Methyl
1-methyl-1H-pyrazol-3-yl
1-methyl-1H-pyrazol-4-yl
1-methyl-1H-pyrazol-5-yl
2-methylphenyl
3-methylphenyl
4-methylphenyl
2-methylpyridin-5-yl
Methylsulfonylmethyl
2-methylsulfonylphenyl
3-methylsulfonylphenyl
4-methylsulfonylphenyl
morpholin-4-ylmethyl
Phenyl
Pyrazinyl
1H-pyrazol-3-yl
pyridin-2-yl
pyridin-3-yl
pyridin-4-yl
3-pyridinylmethyl
pyrimidin-2-yl
pyrimidin-4-yl
pyrimidin-5-yl
quinolin-4-yl
quinolin-5-yl
quinolin-8-yl
5,6,7,8-tetrahydro-1,8-naphthyridin-2-yl

6,7,8,9-tetrahydro-5H-benzo[a][7]annulen-5-yl
6,7,8,9-tetrahydro-5H-benzo[a][7]annulen-6-yl
Tetrahydrofuran-2-yl
1,2,3,4-tetrahydronaphthalen-1-yl
1,2,3,4-tetrahydronaphthalen-2-yl
1,3-thiazol-2-yl
1,3-thiazol-5-yl
thiophen-2-yl
thiophen-3-yl

20. The compound according to claim 1, selected from the group consisting of:

- 5 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{2-[3-(methylsulfonyl)phenyl]-4-phenyl-1,3-thiazol-5-yl}phenyl)quinoline,
- 2-[4-(3-chlorophenyl)-5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-1,3-thiazol-2-yl]propan-2-ol,
- 2-[4-(4-fluorophenyl)-5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-1,3-thiazol-2-yl]propan-2-ol,
- 10 8-{3-[4-(4-chlorophenyl)-2-quinolin-5-yl]-1,3-thiazol-5-yl}phenyl}-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline,
- 2-{3-[4-(3-chlorophenyl)-5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-1,3-thiazol-2-yl]phenyl}propan-2-ol,
- 15 2-{3-[4-(3-chloro-4-fluorophenyl)-5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-1,3-thiazol-2-yl]phenyl}propan-2-ol,
- 2-{3-[4-[3,4-bis(difluoromethoxy)phenyl]-5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-1,3-thiazol-2-yl]phenyl}propan-2-ol,
- N*-{5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-2-yl}acetamide,
- 20 *N*-{5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-2-yl}pyridin-4-amine,
- 2-[5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-4-pyridin-4-yl]-1,3-thiazol-2-yl]propan-2-ol,

MC054PV

- 2-[5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-4-(1-oxidopyridin-4-yl)-1,3-thiazol-2-yl]propan-2-ol,  
 2-[5-(4-chlorophenyl)-4-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-1,3-thiazol-2-yl]propan-2-ol,  
 5 2-{3-[4-(4-chlorophenyl)-5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-1,3-thiazol-2-yl]phenyl}propan-2-ol, and  
 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-{3-[5-(1*H*-tetraazol-5-yl)pyridin-3-yl]phenyl}quinoline,  
 or a pharmaceutically acceptable salt thereof.

10

21. The compound according to claim 1, selected from the group consisting of:
- 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{5-[4-(methylsulfonyl)phenyl]-1-oxidopyridin-3-yl}phenyl)quinoline,  
 15 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{5-[3-(methylsulfonyl)phenyl]-1-oxidopyridin-3-yl}phenyl)quinoline,  
 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{5-[2-(methylsulfonyl)phenyl]-1-oxidopyridin-3-yl}phenyl)quinoline,  
 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{2-[4-(methylsulfonyl)phenyl]pyridin-3-yl}phenyl)quinoline,  
 20 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-[3-(1-oxido-5-phenylpyridin-3-yl)phenyl]quinoline,  
 8-{3-[5-(3,5-dichlorophenyl)-1-oxidopyridin-3-yl]phenyl}-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline,  
 25 8-{3-[5-(3,4-dimethoxyphenyl)-1-oxidopyridin-3-yl]phenyl}-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline,  
 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-{3-[5-(5-methyl-1,2,4-oxadiazol-3-yl)-1-oxidopyridin-3-yl]phenyl}quinoline,  
 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-{3-[5-(5-methyl-1,3,4-oxadiazol-2-yl)-1-oxidopyridin-3-yl]phenyl}quinoline,  
 30 8-{3-[6-(benzyloxy)-5-(3-methyl-1,2,4-oxadiazol-5-yl)pyridin-3-yl]phenyl}-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline,  
 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{6-[4-(methylsulfonyl)phenyl]-1-oxidopyridin-3-yl}phenyl)quinoline,

- 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{5-[4-(methylsulfonyl)phenyl]-1-oxidopyridin-2-yl}phenyl)quinoline,  
 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-{3-[3-[4-(methylsulfonyl)phenyl]-5-(trifluoromethyl)pyridin-2-yl]phenyl}quinoline,  
 5 1-(4-chlorophenyl)-3-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)pyridin-2(1*H*)-one,  
*N*-isopropyl-5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-4-[4-(methylsulfonyl)phenyl]thiophene-2-carboxamide,  
 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(1,1':2',1''-terphenyl-3-yl)quinoline,  
 10 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-[4'-(methylthio)-1,1':2',1''-terphenyl-3-yl]quinoline,  
 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-[2'-(5-methyl-1,3,4-oxadiazol-2-yl)-1,1'-biphenyl-3-yl]quinoline,  
 methyl 3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}-4''-(methylsulfonyl)-  
 15 1,1':2',1''-terphenyl-4'-carboxylate,  
 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(1,1':4',1''-terphenyl-3-yl)quinoline,  
 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(1,1':3',1''-terphenyl-3-yl)quinoline,  
 2-[5-(3'-(6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl)-1,1'-biphenyl-2-yl)-1,3-thiazol-2-yl]propan-2-ol,  
 20 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-[3'-(1-oxidopyridin-4-yl)-1,1'-biphenyl-3-yl]quinoline,  
 or a pharmaceutically acceptable salt thereof.
22. The compound according to claim 1, selected from the group consisting of:
- 25 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{4-[4-(methylsulfonyl)phenyl]pyridin-3-yl}phenyl)quinoline,  
 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{4-[4-(methylsulfonyl)phenyl]-1-oxidopyridin-3-yl}phenyl)quinoline,  
 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{4-[3-(methylthio)phenyl]pyridin-3-yl}phenyl)quinoline,  
 30 8-[4',5'-difluoro-4''-(methylthio)-1,1':2',1''-terphenyl-3-yl]-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline,  
 8-[4',5'-difluoro-4''-(methylsulfonyl)-1,1':2',1''-terphenyl-3-yl]-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline,

MC054PV

- 8-(4"-fluoro-1,1':2',1"-terphenyl-3-yl)-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline,  
 6,7-dichloro-2-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-3-[4-(methylsulfonyl)phenyl]quinoxaline,  
 2-(4-chlorophenyl)-3-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)quinoxaline,  
 2-{4-[3-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)quinoxalin-2-yl]phenyl}propan-2-ol,  
 2-[3,4-bis(difluoromethoxy)phenyl]-3-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)quinoxaline,  
 4-[3-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)quinoxalin-2-yl]benzoic acid,  
*N*-cyclopropyl-4-[3-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)quinoxalin-2-yl]benzamide,  
 2-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-3-(4-methylphenyl)quinoxaline,  
 2-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-3-phenylquinoxaline,  
 2-(4-fluorophenyl)-3-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)quinoxaline,  
 2-{4-[3-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)pyrazin-2-yl]phenyl}propan-2-ol,  
 6-[1-methyl-1-(methylsulfonyl)ethyl]-8-(3-{3-[4-(methylthio)phenyl]pyrazin-2-yl}phenyl)quinoline,  
 8-{3-[3-(4-fluorophenyl)pyrazin-2-yl]phenyl}-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline,  
 8-(3-{2-(2-ethylpyridin-4-yl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-5-yl}phenyl)-6-[1-methyl-1-(methylsulfonyl)ethyl]quinoline,  
 2-(4-{5-(3-{6-[1-methyl-1-(methylsulfonyl)ethyl]quinolin-8-yl}phenyl)-4-[4-(methylsulfonyl)phenyl]-1,3-thiazol-2-yl}phenyl)propan-2-ol,  
 or a pharmaceutically acceptable salt thereof.

23. A pharmaceutical composition comprising a therapeutically effective amount of the compound according to claim 1 or a pharmaceutically acceptable salt thereof; and

MC054PV

a pharmaceutically acceptable carrier.

24. The pharmaceutical composition according to claim 18, further comprising a Leukotriene receptor antagonist, a Leukotriene biosynthesis inhibitor, an M2/M3 antagonist, a corticosteroid, an H1 receptor antagonist or a beta 2 adrenoceptor agonist.

25. The pharmaceutical composition according to claim 18, further comprising a COX-2 selective inhibitor, a statin, or an NSAID.

26. A method of treatment or prevention of asthma; chronic bronchitis; chronic obstructive pulmonary disease; adult respiratory distress syndrome; infant respiratory distress syndrome; cough; chronic obstructive pulmonary disease in animals; adult respiratory distress syndrome; ulcerative colitis; Crohn's disease; hypersecretion of gastric acid; bacterial, fungal or viral induced sepsis or septic shock; endotoxic shock; laminitis or colic in horses; spinal cord trauma; head injury; neurogenic inflammation; pain; reperfusion injury of the brain; psoriatic arthritis; rheumatoid arthritis; ankylosing spondylitis; osteoarthritis; inflammation; or cytokine-mediated chronic tissue degeneration comprising the step of administering a therapeutically effective amount, or a prophylactically effective amount, of the compound according to claim 1 or a pharmaceutically acceptable salt thereof.

27. A method of treatment or prevention of allergic rhinitis, allergic conjunctivitis, eosinophilic granuloma, osteoporosis, arterial restenosis, atherosclerosis, reperfusion injury of the myocardium chronic glomerulonephritis, vernal conjunctivitis, cachexia, transplant rejection, or graft versus host disease, comprising the step of administering a therapeutically effective amount, or a prophylactically effective amount, of the compound according to claim 1 or a pharmaceutically acceptable salt thereof.

28. A method of treatment or prevention of depression, memory impairment, monopolar depression, Parkinson disease, Alzheimer's disease, acute and chronic multiple sclerosis, psoriasis, benign or malignant proliferative skin diseases, atopic dermatitis, urticaria, cancer, tumor growth or cancerous invasion of normal tissues, comprising the step of administering a therapeutically effective amount, or a

MC054PV

prophylactically effective amount, of the compound according to claim 1 or a pharmaceutically acceptable salt thereof.

5

MC054PV

## TITLE OF THE INVENTION

8-(3-BIARYL)PHENYLQUINOLINE PHOSPHODIESTERASE-4 INHIBITORS

## ABSTRACT OF THE DISCLOSURE

5 Novel substituted 8-phenylquinolines, wherein the phenyl group at the 8-position contains an aryl or heteroaryl substituent in the *meta* position, are PDE4 inhibitors.



**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**